



Nonprofit corporations and independent licensees  
of the Blue Cross and Blue Shield Association

Medical benefit drug policies are a source for BCBSM and BCN medical policy information only. These documents are not to be used to determine benefits or reimbursement. Please reference the appropriate certificate or contract for benefit information. This policy may be updated and therefore subject to change.

**Effective Date: 12/14/2023**

**Ilumya™ (tildrakizumab-asmn)**

**HCPCS: J3245**

**Policy:**

*Requests must be supported by submission of chart notes and patient specific documentation.*

- A. Coverage of the requested drug is provided when all the following are met:
  - a. FDA approved age
  - b. Diagnosis of psoriasis (PsO)
    - i. Trial and failure, contraindication, or intolerance to one topical corticosteroid
  - c. Not to be used in combination with other biologics or targeted disease-modifying anti-rheumatic drugs (DMARDs)
  - d. Trial and failure, contraindication, OR intolerance to the preferred drugs as listed in BCBSM/BCN's utilization management medical drug list.
  
- B. Quantity Limitations, Authorization Period and Renewal Criteria
  - a. Quantity Limits: Align with FDA recommended dosing
  - b. Authorization Period: One year at a time
  - c. Renewal Criteria: Clinical documentation must be provided to confirm that current criteria are met and that the medication is providing clinical benefit

\*\*\*Note: Coverage and approval duration may differ for Medicare Part B members based on any applicable criteria outlined in Local Coverage Determinations (LCD) or National Coverage Determinations (NCD) as determined by Center for Medicare and Medicaid Services (CMS). See the CMS website at <http://www.cms.hhs.gov/>. Determination of coverage of Part B drugs is based on medically accepted indications which have supported citations included or approved for inclusion determined by CMS approved compendia.

**Background Information:**

- Ilumya is a an interleukin-23 (IL-23) antagonist indicated for the treatment of moderate-to-severe psoriasis in adults who are candidates for systemic therapy or phototherapy. Per the prescribing information, Ilumya should only be administered by a healthcare provider.
  
- Psoriasis is a chronic, painful and life-altering immune-mediated disease which predominantly manifests with skin and joint involvement. Patients may also experience significant cardiovascular and psychological comorbidities.

This policy and any information contained herein is the property of Blue Cross Blue Shield of Michigan and its subsidiaries, is strictly confidential, and its use is intended for the P&T committee, its members and BCBSM employees for the purpose of coverage determinations.

Approximately 2% of U.S. adults are affected by psoriasis (men and women equally), and it can occur at any age. Approximately 90% of psoriasis-affected patients have plaque psoriasis, which is characterized by well-defined round or oval plaques that vary in size and often coalesce. The severity of psoriasis is defined as: mild = less than 3% of body affected; moderate = 3-10% of body affected; and severe being more than 10% of the body affected.

- Per the 2020 Joint American Academy of Dermatology - National Psoriasis Foundation (AAD/NPF) guidelines of care for the management and treatment of psoriasis with topical therapy and alternative medicine modalities for psoriasis severity measures: topical corticosteroids provide a high efficacy and good safety option for patients with localized disease. They are generally recommended as first-line therapy. Choice of steroid potency may depend on severity, location, patient preference, and patient age, while the duration of treatment may vary with steroid potency, location and severity of disease often ranging from 2-12 weeks. Therapeutic regimens may include 2-4 weeks with a topical steroid applied twice daily, followed by a maintenance regimen where topical steroids are alternated with a steroid-sparing topical agent. Treatment with topical steroids for over 12 weeks is recommended under careful supervision by a physician.
- Per the 2019 Joint AAD/NPF guidelines of care for the management and treatment of psoriasis with phototherapy: phototherapy serves as a reasonable and effective treatment option for patients requiring more than topical medications and/or those wishing to avoid systemic medications or simply seeking an adjunct to a failing regimen. Guidelines also state that the majority of patients with mild-to-moderate disease have adequate disease control with topical therapies and phototherapy alone.
- Per the Joint AAD/NPF guidelines of care for the management and treatment of psoriasis with systemic nonbiologic therapies: many oral medications, including methotrexate, cyclosporine, and acitretin, have been used for decades to treat psoriasis, each with its own benefits and risks. Most work by targeting the immune system, whereas others, such as acitretin, work predominantly by decreasing keratinocyte hyperproliferation, thus restoring the normal epidermal differentiation. Both methotrexate and cyclosporine are category A guideline recommendations for the treatment of moderate to severe psoriasis in adults and for severe, recalcitrant psoriasis, respectively. Studies examining the use of methotrexate and cyclosporine in psoriasis showed the primary efficacy endpoints met within 12-16 weeks. Acitretin is a category B guideline recommendation as monotherapy for plaque psoriasis, with full treatment response expected within 3-6 months.
- Per the 2019 Joint AAD/NPF guidelines of care for the management and treatment of psoriasis with biologics: biologic agents, as monotherapy or combined with other topical or systemic medications, have a high benefit-to-risk ration. Tumor necrosis factor (TNF)-alpha and IL-12/IL-23, IL-23, and IL-17 products have a category "A" recommendation as a monotherapy treatment option for adult patients with moderate-to-severe plaque psoriasis. Guidelines do not recommend one product over another and note the similar efficacy seen across biologics within the same class. There are no published, robust studies to support the use of more than one biologic product or targeted DMARD in combination.

## References:

1. Ilumya [prescribing information]. Whitehouse Station, NJ, Merck & Co., Inc., March 2018.
2. Elmets CA, Lim HW, Stoff B, et al. Joint American Academy of Dermatology-National Psoriasis Foundation guidelines of care for the management and treatment of psoriasis with phototherapy [published correction appears in J Am Acad Dermatol. 2020 Mar;82(3):780]. J Am Acad Dermatol. 2019;81(3):775-804. doi:10.1016/j.jaad.2019.04.042
3. Elmets CA, Korman NJ, Prater EF, et al. Joint AAD-NPF Guidelines of care for the management and treatment of psoriasis with topical therapy and alternative medicine modalities for psoriasis severity measures [published online ahead of print, 2020 Jul 30]. J Am Acad Dermatol. 2020;S0190-9622(20)32288-X. doi:10.1016/j.jaad.2020.07.087

This policy and any information contained herein is the property of Blue Cross Blue Shield of Michigan and its subsidiaries, is strictly confidential, and its use is intended for the P&T committee, its members and BCBSM employees for the purpose of coverage determinations.

4. Menter A, Gelfand JM, Connor C, et al. Joint American Academy of Dermatology-National Psoriasis Foundation guidelines of care for the management of psoriasis with systemic nonbiologic therapies. J Am Acad Dermatol. 2020;82(6):1445-1486. doi:10.1016/j.jaad.2020.02.044
5. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. J Am Acad Dermatol. 2019;80(4):1029-1072. doi:10.1016/j.jaad.2018.11.057

Policy History												
#	Date	Change Description										
1.9	Effective Date: 12/14/2023	Annual review of criteria performed, no changes were made										
1.8	Effective Date: 12/01/2022	Annual review of criteria performed, no changes were made										
1.7	Effective Date: 02/04/2021	Removed FDA approved indications criteria, phototherapy and oral DMARD criteria for psoriasis, and added trial of one topical corticosteroid for psoriasis										
1.6	Effective Date: 02/04/2021	Removal of the topical steroid criteria for psoriasis indication										
1.5	Effective Date: 12/03/2020	Previously approved criteria updated to align management between pharmacy and medical benefit for all listed indications										
1.4	Effective Date: 08/13/2020	Annual Review of Medical Policy										
1.3	Effective Date: 08/15/2019	Annual Review of Medical Policy										
1.2	Effective Date: 08/09/2018	New Drug Review										
1.1	Effective Date: 08/01/2018	UM medical management system update for BCBS and BCN <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>Line of Business</th> <th>PA Required in Medical Management System (Yes/No)</th> </tr> </thead> <tbody> <tr> <td>BCBS</td> <td>Yes</td> </tr> <tr> <td>BCN</td> <td>Yes</td> </tr> <tr> <td>MAPPO</td> <td>No</td> </tr> <tr> <td>BCNA</td> <td>No</td> </tr> </tbody> </table>	Line of Business	PA Required in Medical Management System (Yes/No)	BCBS	Yes	BCN	Yes	MAPPO	No	BCNA	No
Line of Business	PA Required in Medical Management System (Yes/No)											
BCBS	Yes											
BCN	Yes											
MAPPO	No											
BCNA	No											
1.0	Effective Date: 05/03/2018	Preliminary Drug Review										

\* The prescribing information for a drug is subject to change. To ensure you are reading the most current information it is advised that you reference the most updated prescribing information by visiting the drug or manufacturer website or <http://dailymed.nlm.nih.gov/dailymed/index.cfm>.