
Medical Policy



Nonprofit corporations and independent licensees
of the Blue Cross and Blue Shield Association

Joint Medical Policies are a source for BCBSM and BCN medical policy information only. These documents are not to be used to determine benefits or reimbursement. Please reference the appropriate certificate or contract for benefit information. This policy may be updated and is therefore subject to change.

***Current Policy Effective Date: 9/1/24**
(See policy history boxes for previous effective dates)

Title: Laboratory Tests Post Transplant (Kidney, Heart, and Lung) and for Heart Failure

Description/Background

Clinical assessment and noninvasive imaging of chronic heart failure can be limited in accurately diagnosing patients with heart failure because symptoms and signs can poorly correlate with objective methods of assessing cardiac dysfunction. For management of heart failure, clinical signs and symptoms (e.g., shortness of breath) are relatively crude markers of decompensation and occur late in the course of an exacerbation. Thus, circulating biomarkers have potential benefit in heart failure diagnosis and management.

In transplant recipients, despite the progress in immunosuppressant therapy, risk of rejection remains. Diagnosis of allograft rejection continues to rely on clinical monitoring and histologic confirmation by tissue biopsy. However, due to limitations of tissue biopsy, including a high degree of interobserver variability in the grading of results and its potential complications, less invasive alternatives have been investigated. Several laboratory-tested biomarkers of transplant rejection have been evaluated and are commercially available for use. The laboratory tests for heart transplant rejection currently evaluated in this policy include the Presage® ST2 Assay kit, which measures the soluble suppression of tumorigenicity-2 (sST2) protein biomarker; the Heartsbreath test, which measures breath markers of oxidative stress; the AlloSure, Prospera Heart, Viracor TRAC, and myTAIHEART tests for assessment of donor-derived cell-free DNA (dd-cfDNA); the AlloMap test, which uses gene expression profiling (GEP); and the HeartCare test, which combines AlloMap GEP testing with the AlloSure test. Also included in this policy are the AlloSure and Prospera dd-cfDNA tests for assessment of renal and lung transplant rejection.

Heart Failure

Heart failure is a major cause of morbidity and mortality worldwide. The term heart failure refers to a complex clinical syndrome that impairs the heart's ability to move blood through the circulatory system.(1) The prevalence of heart failure in the U.S. between 2013 and 2016 was an estimated 6.2 million for Americans ≥ 20 years old, up from 5.7 million from between 2009 and 2012.(2,3) Heart failure is the leading cause of hospitalization among people older than age 65 years, with direct and indirect costs estimated at \$37 billion annually in the U.S.(2) Although survival has improved with treatment advances, absolute mortality rates of heart failure remain near 50% within 5 years of diagnosis.

Physiology

Heart failure can be caused by disorders of the pericardium, myocardium, endocardium, heart valves or great vessels, or metabolic abnormalities. Individuals with heart failure may present with a wide range of left ventricular (LV) anatomy and function. Some have normal LV size and preserved ejection fraction; others have severe LV dilatation and depressed ejection fraction. However, most patients present with key signs and symptoms secondary to congestion in the lungs from impaired LV myocardial function.(1) They include dyspnea, orthopnea, and paroxysmal dyspnea. Other symptoms include weight gain due to fluid retention, fatigue, weakness, and exercise intolerance secondary to diminished cardiac output.

Diagnosis

Initial evaluation of a patient with suspected heart failure is typically based on clinical history, physical examination, and chest radiograph. Because people with heart failure may present with nonspecific signs and symptoms (e.g., dyspnea), accurate diagnosis can be challenging. Therefore, noninvasive imaging procedures (e.g., echocardiography, radionuclide angiography) are used to quantify pump function of the heart, thus identifying or excluding heart failure in patients with characteristic signs and symptoms. These tests can also be used to assess prognosis by determining the severity of the underlying cardiac dysfunction.(1) However, clinical assessment and noninvasive imaging can be limited in accurately evaluating patients with heart failure because symptoms and signs can poorly correlate with objective methods of assessing cardiac dysfunction.(4,5,6) Thus, invasive procedures (e.g., cardiac angiography, catheterization) are used in select patients with presumed heart failure symptoms to determine the etiology (i.e., ischemic vs. nonischemic) and physiologic characteristics of the condition.

Treatment

Patients with heart failure may be treated using a number of interventions. Lifestyle factors such as the restriction of salt and fluid intake, monitoring for increased weight, and structured exercise programs are beneficial components of self-management. A variety of medications are available to treat heart failure. They include diuretics (e.g., furosemide, hydrochlorothiazide, spironolactone), angiotensin-converting enzyme inhibitors (e.g., captopril, enalapril, lisinopril), angiotensin receptor blockers (e.g., losartan, valsartan, candesartan), b-blockers (e.g., carvedilol, metoprolol succinate), and vasodilators (e.g., hydralazine, isosorbide dinitrate). Numerous device-based therapies also are available. Implantable cardioverter defibrillators reduce mortality in patients with an increased risk of sudden cardiac death. Cardiac resynchronization therapy improves symptoms and reduces mortality for patients who have disordered LV conduction evidenced by a wide QRS complex on electrocardiogram. Ventricular assist devices are indicated for patients with end stage heart failure who have failed all other therapies and are also used as a bridge to cardiac transplantation in select patients.(1)

Heart Failure Biomarkers

Because of limitations inherent in standard clinical assessments of patients with heart failure, a number of objective disease biomarkers have been investigated to diagnose and assess heart failure patient prognosis, with the additional goal of using biomarkers to guide therapy.(7) They include a number of proteins, peptides, or other small molecules whose production and release into circulation reflect the activation of remodeling and neurohormonal pathways that lead to LV impairment. Examples include B-type natriuretic peptide (BNP), its analogue N-terminal pro B-type natriuretic peptide (NT-proBNP), troponin T and I, renin, angiotensin, arginine vasopressin, C-reactive protein, and norepinephrine.(1,7)

BNP and NT-proBNP are considered the reference standards for biomarkers in assessing heart failure patients. They have had substantial impact on the standard of care for diagnosis of heart failure and are included in the recommendations of all major medical societies, including the American College of Cardiology Foundation and American Heart Association,(8) European Society of Cardiology,(9) and the Heart Failure Society of America.(10) Although natriuretic peptide levels are not 100% specific for the clinical diagnosis of heart failure, elevated BNP or NT-proBNP levels in the presence of clinical signs and symptoms reliably identify the presence of structural heart disease due to remodeling and heightened risk for adverse events. Natriuretic peptides also can help in determining prognosis of heart failure patients, with elevated blood levels portending poorer prognosis.

In addition to diagnosing and assessing prognosis of heart failure patients, blood levels of BNP or NT-proBNP have been proposed as an aid for managing patients diagnosed with chronic heart failure.(8,10,12) Levels of either biomarker rise in response to myocardial damage and LV remodeling, whereas they tend to fall as drug therapy ameliorates symptoms of heart failure. Evidence from a large number of randomized controlled trials (RCTs) that have compared BNP- or NTproBNP-guided therapy with clinically guided adjustment of pharmacologic treatment of patients who had chronic heart failure has been assessed in recent systematic reviews and meta-analyses. However, these analyses have not consistently reported a benefit for BNP-guided management. Savarese et al (2013) published the largest meta-analysis to date, a patient-level meta-analysis that evaluated 2686 patients from 12 RCTs.(11) This meta-analysis showed that NTproBNP- guided management was associated with significant reductions in all-cause mortality and heart failure–related hospitalization compared with clinically guided treatment. Although BNP-guided management in this meta-analysis was not associated with significant reductions in these parameters, differences in patient numbers and characteristics may explain the discrepancy. Troughton et al (2014) conducted a second patient-level meta-analysis that included 11 RCTs with 2000 patients randomized to natriuretic peptide-guided pharmacologic therapy or usual care.(12) The results showed that, among patients 75 years of age or younger with chronic heart failure, most of whom had impaired left ventricular ejection fraction, natriuretic peptide-guided therapy was associated with significant reductions in all-cause mortality compared with clinically guided therapy. Natriuretic-guided therapy also was associated with significant reductions in hospitalization due to heart failure or cardiovascular disease.

Suppression of Tumorigenicity-2 Protein Biomarker

A protein biomarker, ST2, has elicited interest as a potential aid to predict prognosis and manage therapy of heart failure.(13-19) This protein is a member of the interleukin-1 (IL-1) receptor family. It is found as a transmembrane isoform (ST2L) and a soluble isoform (sST2), both of which have circulating IL-33 as their primary ligand. ST2 is a unique biomarker that has pluripotent effects in vivo. Thus, binding between IL-33 and ST2L is believed to have an immunomodulatory function via T-helper type 2 lymphocytes and was initially described in the context of cell proliferation, inflammatory states, and autoimmune diseases.(20) However, the IL-33/ST2L signaling cascade is also strongly induced through mechanical strain of cardiac fibroblasts or cardiomyocytes. The net result is mitigation of adverse cardiac remodeling and myocardial fibrosis, which are key processes in the development of heart failure.(21) The soluble isoform of ST2 is produced by lung epithelial cells and cardiomyocytes and is secreted into circulation in response to exogenous stimuli, mechanical stress, and cellular stretch. This form of ST2 binds to circulating IL-33, acting as a "decoy," thus inhibiting the IL-33-associated anti-remodeling effects of the IL-33/ST2L signaling pathway. Thus, on a biologic level, IL-33/ST2L signaling plays a role in modulating the balance of inflammation and neurohormonal activation and is viewed as pivotal for protection from myocardial remodeling, whereas sST2 is viewed as attenuating this protection. In the clinic, blood concentrations of sST2 appear to correlate closely with adverse cardiac structure and functional changes consistent with remodeling in patients with heart failure, including abnormalities in filling pressures, chamber size, and systolic and diastolic function.(7,15,17)

An enzyme-linked immunosorbent-based assay is commercially available for determining sST2 blood levels (Presage ST2 Assay).(18) The manufacturer claims a limit of detection of 1.8 ng/mL for sST2, and a limit of quantification of 2.4 ng/mL, as determined according to Clinical and Laboratory Standards Institute guideline EP-17-A.Mueller and Dieplinger (2013) reported a limit of detection of 2.0 ng/mL for sST2 in their study.(18) In the same study, the assay had a within-run coefficient of variation of 2.5% and a total coefficient of variation less than 4.0%, demonstrated linearity within the dynamic range of the assay calibration curve, and exhibited no relevant interference or cross-reactivity.

The ST2 biomarker is not intended to diagnosis heart failure because it is a relatively nonspecific marker that is increased in many other disparate conditions that may be associated with acute or chronic manifestations of heart failure.(17,18) Although the natriuretic peptides (BNP, NT-proBNP) reflect different physiologic aspects of heart failure compared with sST2, they are considered the reference standard biomarkers when used with clinical findings to diagnose, prognosticate, and manage heart failure and as such are the comparator to sST2.

HEART TRANSPLANT REJECTION

Most cardiac transplant recipients experience at least a single episode of rejection in the first year after transplantation. The International Society for Heart and Lung Transplantation (2005) modified its grading scheme for categorizing cardiac allograft rejection.(22) The revised (R) categories are listed in Table 1.

Table 1. Revised Grading Schema for Cardiac Allograft Rejection

New Grade	Definition	Old Grade
0R	No rejection	
1R	Mild rejection	1A, 1B, and 2
2R	Moderate rejection	3A
3R	Severe rejection	3B and 4

Acute cellular rejection is most likely to occur in the first 6 months after transplantation, with a significant decline in the incidence of rejection after this time. Although immunosuppressants are required on a life-long basis, dosing is adjusted based on graft function and the grade of acute cellular rejection determined by histopathology. Endomyocardial biopsies are typically taken from the right ventricle via the jugular vein periodically during the first 6 to 12 months posttransplant. The interval between biopsies varies among clinical centers. A typical schedule is weekly for the first month, once or twice monthly for the following 6 months, and several times (monthly to quarterly) between 6 months and 1-year posttransplant. Surveillance biopsies may also be performed after the first postoperative year (e.g., on a quarterly or semiannual basis). This practice, although common, has not been demonstrated to improve transplant outcomes. Some centers no longer routinely perform endomyocardial biopsies after 1 year in patients who are clinically stable.

While the endomyocardial biopsy is the criterion standard for assessing heart transplant rejection, it is limited by a high degree of interobserver variability in the grading of results and potential morbidity that can occur with the biopsy procedure. Also, the severity of rejection may not always coincide with the grading of the rejection by biopsy. Finally, a biopsy cannot be used to identify patients at risk of rejection, limiting the ability to initiate therapy to interrupt the development of rejection. For these reasons, an endomyocardial biopsy is considered a flawed criterion standard by many. Therefore, noninvasive methods of detecting cellular rejection have been explored. It is hoped that noninvasive tests will assist in determining appropriate patient management and avoid overuse or underuse of treatment with steroids and other immunosuppressants that can occur with false-negative and false-positive biopsy reports. Two techniques are commercially available for the detection of heart transplant rejection.

Noninvasive Heart Transplant Rejection Tests

Presage ST2 Assay

In addition to its use as a potential aid to predict prognosis and manage therapy of heart failure, elevated serum ST2 levels have also been associated with an increased risk of antibody-mediated rejection following a heart transplant. For this reason, ST2 has also been proposed as a prognostic marker post heart transplantation and as a test to predict acute cellular rejection (graft-versus-host disease). The Presage ST2 Assay, described above, is a commercially available sST2 test that has been investigated as a biomarker of heart transplant rejection.

Heartsbreath Test

The Heartsbreath test, a noninvasive test that measures breath markers of oxidative stress, has been developed to assist in the detection of heart transplant rejection. In heart transplant recipients, oxidative stress appears to accompany allograft rejection, which degrades membrane polyunsaturated fatty acids and evolving alkanes and methylalkanes that are in turn, excreted as volatile organic compounds in breath. The Heartsbreath test analyzes the breath methylated alkane contour, which is derived from the abundance of C4 to C20 alkanes and monomethylalkanes and has been identified as a marker to detect grade 3 (clinically significant) heart transplant rejection.

HeartCare

Cell-free DNA (cfDNA), released by damaged cells, is normally present in healthy individuals.(23) In patients who have received transplants, donor-derived cell-free DNA (dd-cfDNA) may be also present. It is proposed that allograft rejection, which is associated with damage to transplanted cells, may result in an increase in dd-cfDNA. HeartCare (CareDx) is a commercially-available test that combines AlloMap gene expression profiling with a next-generation sequencing assay that quantifies the fraction of dd-cfDNA in cardiac transplant recipients relative to total cfDNA. The AlloMap score, AlloMap score variability, and AlloSure % dd-cfDNA are reported.

Prospera

Prospera Heart (Natera) is a commercially available assay that uses massively multiplexed PCR (mmPCR) followed by next-generation sequencing (NGS) to quantify the fraction of dd-cfDNA in transplant recipients. Donor versus recipient cfDNA is differentiated via an advanced bioinformatics analysis of >13,000 single-nucleotide polymorphisms (SNPs) without the need for prior recipient or donor genotyping or computational adjustments for related donors.(24) The Prospera Heart test reports the dd-cfDNA fraction in the patient's blood as a predictor of acute rejection, although the optimal dd-cfDNA cut-point is not described by the manufacturer.

myTAIHEART

Using proprietary myTAIHEART software (TAI Diagnostics), the myTAIHEART test uses multiplexed, high-fidelity amplification followed by allele-specific qPCR of a panel of 94 highly informative bi-allelic single nucleotide polymorphisms (SNPs) and two controls to quantitatively genotype cfDNA in the patient's plasma after cardiac transplant, and accurately distinguish dd-cfDNA originating from the engrafted heart from cfDNA originating from the recipient's native cells.(25) The ratio of dd-cfDNA to total cfDNA is reported as the donor fraction (%) and categorizes the patient as at low or increased risk of moderate (grade 2R) to severe (grade 3R) acute cellular rejection: low donor fractions indicate less damage to the transplanted heart and a lower risk for rejection, while increased donor fractions indicate more damage to the transplanted heart and an increased risk for rejection. Testing with myTAIHEART does not require a donor specimen. TAI Diagnostics suspended production of the myTAIHEART test in 2020. As of September 2022, TAI Diagnostics appears to no longer be operational and it is unclear if myTAIHEART will be available through another company in the future.

AlloMap

Another approach has focused on patterns of gene expression of immunomodulatory cells, as detected in the peripheral blood. For example, microarray technology permits the analysis of the expression of thousands of genes, including those with functions known or unknown. Patterns of gene expression can then be correlated with known clinical conditions, permitting a selection of a finite number of genes to compose a custom multigene test panel, which then can be evaluated using polymerase chain reaction techniques. AlloMap (CareDx) is a commercially available molecular expression test that has been developed to detect acute heart transplant rejection or the development of graft dysfunction. The test involves polymerase chain reaction-expression measurement of a panel of genes derived from peripheral blood cells and applies an algorithm to the results. The proprietary algorithm produces a single score that considers the contribution of each gene in the panel. The score ranges from 0 to 40. The AlloMap website states that a lower score indicates a lower risk of graft rejection; the website does not cite a specific cutoff for a positive test.(26) All AlloMap testing is performed at the CareDx reference laboratory in California.

Other laboratory-tested biomarkers of heart transplant rejection have been evaluated. They include brain natriuretic peptide, troponin, and soluble inflammatory cytokines. Most have had low accuracy in diagnosing rejection. Preliminary studies have evaluated the association between heart transplant rejection and micro-RNAs or high-sensitivity cardiac troponin in cross-sectional analyses but the clinical use has not been evaluated.(27,28)

RENAL TRANSPLANT REJECTION

Allograft dysfunction is typically asymptomatic and has a broad differential, including graft rejection. Diagnosis and rapid treatment are recommended to preserve graft function and prevent loss of the transplanted organ. For a primary kidney transplant from a deceased donor (accounting for about 70% of kidney donors), graft survival at 1 year is 95%; at 5 years, graft survival is 78%.(29, 30)

Surveillance of transplant kidney function relies on routine monitoring of serum creatinine, urine protein levels, and urinalysis.(31) Allograft dysfunction may also be demonstrated by a drop in urine output or, rarely, as pain over the transplant site. With clinical suspicion of allograft dysfunction, additional noninvasive workup including ultrasonography or radionuclide imaging may be used. A renal biopsy allows a definitive assessment of graft dysfunction and is typically a percutaneous procedure performed with ultrasonography or computed tomography guidance. Biopsy of a transplanted kidney is associated with fewer complications than biopsy of a native kidney because the allograft is typically transplanted more superficially than a native kidney. Renal biopsy is a low-risk invasive procedure that may result in bleeding complications; loss of a renal transplant, as a complication of renal biopsy, is rare.(32)

Kidney biopsies allow for diagnosis of acute and chronic graft rejection, which may be graded using the Banff Classification.(33,34) Pathologic assessment of biopsies demonstrating acute rejection allows clinicians to further distinguish between acute cellular rejection and antibody-mediated rejection, which are treated differently.

NONINVASIVE RENAL TRANSPLANT REJECTION TESTS

AlloSure

AlloSure Kidney (CareDx) is a commercially available, next-generation sequencing (NGS) assay which quantifies the fraction of dd-cfDNA in renal transplant recipients, relative to total cfDNA, by measuring 266 single nucleotide variants (SNVs). Separate genotyping of the donor or recipient is not required, but patients who received a kidney transplant from a monozygotic (identical) twin are not eligible for this test. The fraction of dd-cfDNA relative to total cfDNA present in the peripheral blood sample is cited in the report. For patients undergoing surveillance, a routine testing schedule is recommended for longitudinal monitoring.

Prospera

Prospera Kidney (Natera) is a commercially available assay that quantifies the fraction of dd-cfDNA in renal transplant recipients. The manufacturer recommends use of the test when there is clinical suspicion of active rejection and for regular surveillance of subclinical rejection in renal transplant recipients.(35) In a surveillance scenario, regular testing is recommended at 1, 2, 3, 4, 6, 9 and 12 months after renal transplant or most recent rejection.(36) Thereafter, the test should be repeated quarterly. The proportion of dd-cfDNA relative to total cfDNA is

reported, with detection of $\geq 1\%$ dd-cfDNA indicating increased risk for active rejection. The percent dd-cfDNA change between tests is also reported.

Lung Transplant Rejection

Despite advances in induction and maintenance immunosuppressive regimens, lung transplant recipients have a median overall survival of 6 years, with more than a third of patients receiving treatment for acute rejection in the first year after transplant.(93,94) Acute cellular rejection, lymphocytic bronchiolitis, and antibody-mediated rejection are all risk factors for subsequent development of chronic lung allograft dysfunction (CLAD). Pathologic grading of acute cellular rejection is based on the histological assessment of perivascular and interstitial mononuclear cell infiltrates. Antibody-mediated rejection may be clinical (symptomatic or asymptomatic allograft dysfunction) or subclinical (normal allograft function). Key diagnostic criteria established via consensus by the International Society for Heart and Lung Transplantation include the presence of antibodies directed toward donor human leukocyte antigens and characteristic lung histology with or without evidence of complement 4d within the graft.(95) The most common phenotype of CLAD is a persistent obstructive decline in lung function known as bronchiolitis obliterans syndrome (BOS), which is graded based on the degree of decrease in FEV1. Approximately 50% of patients develop BOS within 5 years post-transplant. Median survival following a diagnosis of BOS is 3-5 years. Acute rejection may present with non-specific physical symptoms or be asymptomatic. However, the role of surveillance bronchoscopy for screening asymptomatic patients for acute rejection is controversial, and performance of surveillance bronchoscopies varies across transplant centers.

Noninvasive Lung Transplant Rejection Tests

AlloSure

AlloSure Lung (CareDx) is a commercially available, NGS assay that quantifies the fraction of dd-cfDNA in lung transplant patients relative to total cfDNA by measuring single nucleotide polymorphisms. The test is intended to provide a direct, noninvasive measure of organ injury in lung transplant patients who are undergoing surveillance. Suggested thresholds for severe injury and quiescence are 1%, 0.85%, and $<0.5\%$, respectively.(96)

Prospera

Prospera Lung (Natera) is a commercially available assay that uses the same methodology as Prospera Heart and Prospera Kidney to quantify the fraction of dd-cfDNA in transplant recipients. The Prospera Lung test reports the dd-cfDNA fraction in the patient's blood as a predictor of acute rejection, chronic rejection, or infection although the optimal dd-cfDNA cut-point for each outcome is not described by the manufacturer.(97)

Regulatory Status

The U.S. Food and Drug Administration (FDA) has cleared multiple biomarker tests for detection of heart and renal allograft rejection. Table 2 provides a summary of the biomarker tests currently included in this policy that have FDA clearance.

Table 2. Select Biomarker Tests for Detection of Heart or Renal Allograft Rejection Cleared by the U.S. Food and Drug Administration

Test	Manufacturer	FDA Clearance Type, Product Number	FDA Clearance Date	Indicated Use
Heartsbreath™	Menssana Research	Humanitarian device exemption, H030004	2004	To aid in diagnosing grade 3 heart transplant rejection in patients who have received heart transplants within the preceding year. The device is intended as an adjunct to, and not as a substitute for, endomyocardial biopsy and is also limited to patients who have had endomyocardial biopsy within the previous month.
AlloMap® Molecular Expression Testing	CareDx, formerly XDx	510(k), k073482	2008	The test is to be used in conjunction with clinical assessment, for aiding in the identification of heart transplant recipients with stable allograft function and a low probability of moderate-to-severe transplant rejection. It is intended for patients at least 15 years old who are at least 2 months posttransplant.
Presage® ST2 Assay kit	Critical Diagnostics	510(k), k093758	2011	For use with clinical evaluation as an aid in assessing the prognosis of patients diagnosed with chronic heart failure

FDA: Food and Drug Administration.

Laboratory Developed Tests

There are also commercially available laboratory-developed biomarker tests for detection of heart and renal allograft rejection. Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments. The AlloSure (CareDx) and Prospera (Natera) dd-cfDNA tests are regulated under the Clinical Laboratory Improvement Amendments standards and all testing is performed at the CareDx reference laboratory.

myTAIHEART is also a laboratory developed test (LDT) developed for clinical diagnostic performance exclusively in the College of American Pathologists (CAP) and Clinical Laboratory Improvement Amendment (CLIA) accredited TAI Diagnostics Clinical Reference Laboratory.(25) This test was developed, and its performance characteristics determined by TAI Diagnostics.

These LDTs have not been cleared or approved by the FDA.

Medical Policy Statement

The safety and effectiveness of peripheral blood gene expression profiling (AlloMap™) for the detection of heart transplant rejection has been established. It may be considered a useful therapeutic option when specified criteria have been met.

The measurement of volatile organic compounds (e.g., breath test/Heartsbreath™) for the evaluation of heart transplant rejection is considered experimental/ investigational. The effectiveness and clinical utility of this test has not been clearly established.

The use of Presage ST2® Assay (1) to evaluate the prognosis of individuals diagnosed with chronic heart failure, (2) to guide management (e.g., pharmacologic, device-based, exercise) of individuals diagnosed with chronic heart failure, OR (3) in the post cardiac transplantation period, including but not limited to predicting prognosis and predicting acute cellular rejection, is considered experimental/investigational. The effectiveness and clinical utility of this test has not been clearly established.

The use of peripheral blood gene expression profile test in combination with peripheral blood measurement of donor-derived cell-free DNA (dd-cfDNA) (e.g., HeartCare) in the management of individuals after heart transplantation, including but not limited to the detection of acute heart transplant rejection or heart transplant graft dysfunction, is considered experimental/investigational. The effectiveness and clinical utility of this test has not been clearly established.

The use of peripheral blood measurement of donor-derived cell-free DNA (e.g., AlloSure™, myTAI_{HEART}®, Prospera®, Viracor TRAC®) in the management of individuals after transplantation (renal, cardiac, or lung) to predict (1) prognosis, (2) acute cellular rejection, or (3) graft dysfunction, is considered experimental/investigational. The effectiveness and clinical utility of this test has not been clearly established.

Inclusionary and Exclusionary Guidelines

Inclusions:

Peripheral blood gene expression profiling (AlloMap) may be appropriate as a screening technique for heart transplant rejection in recipients who meet **ALL** of the following:

- At least 15 years old
- 6 months post-heart transplant **AND**

Recipient must have stable heart allograft function demonstrated by **ALL** of the following:

- Left ventricular ejection fraction $\geq 45\%$ which has been confirmed by echocardiogram
- No evidence of CHF
- No evidence of severe cardiac allograft vasculopathy **AND**

Recipient must have a low probability of moderate or severe acute cellular rejection as demonstrated by the **ALL** of the following:

- Clinical assessment (e.g., International Society for Heart and Lung Transplantation rejection status Grade of 0R or 1R)
- No history or evidence of antibody mediated rejection

Exclusions:

- Gene expression profiling (i.e., AlloMap) for any indication not listed above.
 - Measurement of volatile organic compounds (e.g., breath test/Heartsbreath™).
 - Presage ST2 Assay in individuals diagnosed with chronic heart failure or in the post cardiac transplantation period.
 - Peripheral blood measurement of donor-derived cell-free DNA (e.g., AlloSure, myTAI_{HEART}, Prospera, Viracor TRAC®) to predict prognosis, acute cellular rejection or graft dysfunction following renal, cardiac or lung transplantation.
 - The use of peripheral blood gene expression profile test in combination with peripheral blood measurement of donor-derived cell-free DNA (dd-cfDNA) (e.g., HeartCare) in the management of individuals after heart transplantation.
-

CPT/HCPCS Level II Codes *(Note: The inclusion of a code in this list is not a guarantee of coverage. Please refer to the medical policy statement to determine the status of a given procedure.)*

Established codes:

81595

Other codes (investigational, not medically necessary, etc.):

0018M	0055U	0118U	0319U	0320U	81479*
83006	84999				

* Donor derived cell-free DNA (ex. Allosure, Prospera); HeartCare (AlloSure and AlloMap combination)

Policy Guidelines

The U.S. Food and Drug Administration has indicated that the Heartsbreath (Menssana Research) test is only for use as an aid in the diagnosis of grade 3 (now known as grade 2R) heart transplant rejection in patients who have received heart transplants within the preceding year and who have had endomyocardial biopsy within the previous month.

Rationale

Evidence reviews assess whether a medical test is clinically useful. A useful test provides information to make a clinical management decision that improves the net health outcome. That is, the balance of benefits and harms is better when the test is used to manage the condition than when another test or no test is used to manage the condition.

The first step in assessing a medical test is to formulate the clinical context and purpose of the test. The test must be technically reliable, clinically valid, and clinically useful for that purpose. Evidence reviews assess the evidence on whether a test is clinically valid and clinically useful. Technical reliability is outside the scope of these reviews, and credible information on technical reliability is available from other sources.

USE OF SOLUBLE SUPPRESSION OF TUMORIGENICITY-2 LEVELS IN CHRONIC HEART FAILURE PATIENTS

Clinical Context and Test Purpose

The purpose of the of the Soluble Suppression of Tumorigenicity-2 (sST2) assay is to determine prognosis and/or to guide management in individuals with chronic heart failure as an alternative to or an improvement on existing tests and clinical assessment.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with chronic heart failure.

Interventions

The test being considered is sST2 assay to determine prognosis and/or to guide management. Elevated sST2 levels are purported to predict higher risk of poor outcomes.

Comparators

Comparators of interest include standard prognostic markers, including B-type natriuretic peptide levels and clinical assessment.

Outcomes

The general outcomes of interest are overall survival (OS), quality of life, and hospitalizations. Follow-up of 6-12 months would be appropriate to assess quality of life outcomes.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Study Selection Criteria

For the evaluation of clinical validity of sST2 testing, methodologically credible studies were selected using the following principles:

- Reported on the accuracy of the marketed version of the technology (including any algorithms used to calculate scores)
- Included a suitable reference standard
- Individual/sample clinical characteristics were described
- Individual selection criteria were described
- Included a validation cohort separate from the development cohort

Review of Evidence

Subanalyses from Randomized Controlled Trials

A number of clinical studies in which sST2 blood levels were determined using the Presage ST2 Assay have reported that there is an association between ST2 levels and adverse outcomes in patients diagnosed with chronic heart failure. A substantial body of biomarker evidence has been reported retrospectively from subsets of patients enrolled in randomized controlled trials (RCTs) of heart failure interventions. These RCTs include the Valsartan Heart Failure Trial (Val-HeFT);⁽³⁷⁾ Heart Failure: A Controlled Trial Investigating Outcomes of Exercise Training (HF-ACTION);⁽³⁸⁾ Controlled Rosuvastatin Multinational Trial in Heart Failure (CORONA)⁽³⁹⁾; and ProBNP Outpatient Tailored Chronic Heart Failure study (PROTECT).⁽⁴⁰⁾ Although patients in these RCTs were well-characterized and generally well-matched between study arms, the trials were neither intended nor designed specifically to evaluate biomarkers as risk predictors. At present, no prospectively gathered evidence is available from an RCT in which sST2 levels were compared with levels of a B-type natriuretic peptide (BNP or N-terminal pro B-type natriuretic peptide [NT-proBNP]) to predict risk for adverse outcomes among well-defined cohorts of patients with diagnosed chronic heart failure. Key results of larger individual studies are summarized in Table 3.

Table 3. Summary of Selected Clinical Studies of sST2 to Predict Outcomes in Chronic Heart Failure Patients

Study	Population	Mean Age, y	Study Description and Biomarkers	Primary Endpoints	Mean FU	Synopsis of Findings
Ky et al (2011) ⁴¹	Ambulatory CHF (N = 1,141, 75% of Penn HF Study population)	56	Retrospective analysis of sST2 and NT-proBNP levels and their incremental usefulness over clinical SHFM	Mortality or cardiac transplant	2.8 y	<ul style="list-style-type: none"> Elevated sST2 levels associated with increased risk (adjusted P=.002) sST2 in plus NT-proBNP levels showed moderate improvement over SHFM in predicting outcomes (P=.017)
Bayes-Genis et al 2012) ⁴²	Ambulatory decompensated HF (N = 891)	70	Retrospective analysis of sST2 and NT-proBNP levels from consecutive series	Mortality	2.8 y	<ul style="list-style-type: none"> Elevated sST2 and NT-proBNP levels provided independent and additive prognostic information for elevated risk of mortality (P<.001)
Broch et al (2012) ⁴³	Ischemic CHF (N = 1,149, 30% of CORONA RCT)	72	Retrospective analysis of sST2, NT-proBNP, and CRP levels	CV mortality, nonfatal myocardial infarction or stroke	2.6 y	<ul style="list-style-type: none"> Elevated sST2 levels independently associated with increased risk for mortality, hospitalization due to HF, or any CV hospitalization (P<.001) sST2 did not provide additive prognostic information vs. NT-proBNP

Felker et al 2013) ⁴⁴	Ambulatory HF (N = 910, 39% of HF-ACTION RCT)	59	Retrospective analysis of sST2 and NT-proBNP levels	Mortality, hospitalization, functional capacity	2.5 y	<ul style="list-style-type: none"> Elevated sST2 levels independently associated with increased risk for mortality, hospitalization due to HF, or any CV hospitalization ($P<.000$) sST2 and NT-proBNP provided independent prognostic information sST2 did not provide additive prognostic information vs. NT-proBNP
Gaggin et al (2013) ⁴⁵	Recently decompensated CHF (n=151, 100% of PROTECT RCT)	63	Retrospective analysis of sST2 and NT-proBNP levels	Composite outcome (worsening HF, hospitalization for HF, clinically significant CV events)	0.8 y	<ul style="list-style-type: none"> Elevated sST2 levels associated with increased risk for adverse CV outcome ($P<.001$) sST2 and NT-proBNP did not provide independent prognostic information
Anand et al (2014) ⁴⁶	CHF (n=1,650, 33% of Val-HeFT RCT)	63	Retrospective analysis of sST2, NT-proBNP, and other biomarker levels	All-cause mortality and composite outcome (mortality, SCD with resuscitation, hospitalization for HF, or administration of IV inotropic or vasodilator drug for ³⁴ h without hospitalization)		<ul style="list-style-type: none"> Elevated sST2 levels independently associated with increased risk of poor outcomes ($P<.001$) Baseline sST2 levels did not provide substantial prognostic information when added to a clinical model that included NT-proBNP levels
Zhang et al 2015) ⁴⁷	De novo HF or decompensated CHF (N = 1161)	58	Prospective analysis of sST2 in hospitalized sample at 1 center in China	All-cause mortality	1 y	<ul style="list-style-type: none"> Elevated sST2 levels independently associated with increased risk of all-cause mortality ($P<.001$) after adjustment for clinical risk factors and NT-proBNP levels
Dupuy et al 2016) ⁴⁸	HF for ≥ 6 mo (N = 178)	75	Prospective analysis of sST2, NT-proBNP, and other biomarker levels in sample from 1 center in France	All-cause mortality and CV mortality	42 mo ^a	<ul style="list-style-type: none"> Elevated sST2 levels independently associated with increased risk for all-cause mortality and CV mortality ($P<.001$) In multivariate analysis, sST2 and CRP significantly associated with all-

CHF: chronic heart failure; CRP: C-reactive protein; CV: cardiovascular; FU: follow-up; HF: heart failure; HF-ACTION: Heart Failure: A Controlled Trial Investigating Outcomes of Exercise Training; IV: intravenous; NT-proBNP: N-terminal pro B-type natriuretic peptide; PROTECT: ProBNP Outpatient Tailored Chronic Heart Failure study; RCT: randomized controlled trial; SCD: sudden cardiac death; SHFM: Seattle Heart Failure Model; sST2: soluble suppression of tumorigenicity-2; Val-HeFT: Valsartan Heart Failure Trial.
a Median.

Meta-Analyses

Aimo et al (2017) pooled findings of studies on the prognostic value of sST2 for chronic heart failure in a meta-analysis.(49) The meta-analysis selected 7 studies, including post hoc analyses of RCTs, and calculated the association between the Presage ST2 Assay and health outcomes. A pooled analysis of 7 studies found that sST2 was a statistically significant predictor of overall mortality (hazard ratio [HR] = 1.75; 95%CI, 1.37-2.22). Moreover, a pooled analysis of 5 studies found that sST2 was a significant predictor of cardiovascular mortality (HR = 1.79; 95% CI, 1.22 to 2.63).

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from RCTs.

No evidence is available from randomized or nonrandomized controlled studies in which outcomes from groups of well-matched patients managed using serial changes in sST2 blood levels were compared with those managed using the reference standard of BNP or NT-proBNP levels.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

No inferences can be drawn about the clinical utility of sST2 levels for chronic heart failure.

Section Summary: Use of Soluble Suppression of Tumorigenicity-2 in Chronic Heart Failure Patients

Several analyses, mostly retrospective, have evaluated whether sST2 levels are associated with disease prognosis, especially mortality outcomes. Studies mainly found that elevated sST2 levels were statistically associated with elevated risk of mortality. A pooled analysis of study results found that sST2 levels significantly predicted overall mortality and cardiovascular mortality. Several studies, however, found that sST2 test results did not provide additional prognostic information compared with BNP or NT-proBNP levels. In general, it appears that elevated sST2 levels predict higher risk of poor outcomes better than lower levels. The available evidence is limited by interstudy inconsistency and differences in patient characteristics, particularly the severity of heart failure, its etiology, duration, and treatment.

Furthermore, most of the evidence was obtained from retrospective analyses of sST2 levels in subsets of larger patient cohorts within RCTs, potentially biasing the findings. The evidence primarily shows associations between elevated sST2 levels and poor outcomes, but does not go beyond that in demonstrating a clinical connection among biomarker status, treatment received, and clinical outcomes.

USE OF SOLUBLE ST2 SUPPRESSION OF TUMORIGENICITY-2 IN POST-HEART TRANSPLANTATION PATIENTS

Clinical Context and Test Purpose

The purpose of sST2 assay is to determine prognosis and/or to predict acute cellular rejection in individuals with heart transplantation an alternative to or an improvement on existing tests.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with heart transplantation.

Interventions

The test being considered is sST2 assay to determine prognosis and/or to predict acute cellular rejection.

Comparators

Comparators of interest include endomyocardial biopsy for predicting acute cellular rejection.

Outcomes

The general outcomes of interest are OS, quality of life, and hospitalizations.

Table 4. Significant Outcomes for Post-Heart Transplantation Patients

Outcomes	Details	Timing
Morbid events	Short-term and long-term events, such as acute cellular rejection, myocardial infarction, and stroke	30 days, 6 months, 1-5 years
Hospitalizations	Inpatient hospital admissions	30 days, 6 months, 1-5 years

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Study Selection Criteria

For the evaluation of clinical validity of sST2 testing, methodologically credible studies were selected using the following principles:

- Reported on the accuracy of the marketed version of the technology (including any algorithms used to calculate scores)
- Included a suitable reference standard
- Individual/sample clinical characteristics were described
- Individual/sample selection criteria were described
- Included a validation cohort separate from the development cohort

Observational Studies

Serum ST2 levels have been proposed as a prognostic marker post heart transplantation and as a test to predict acute cellular rejection (graft-versus-host disease). There is very little evidence available for these indications. Januzzi et al (2013) retrospectively assessed sST2 levels in 241 patients post-heart transplant.⁽⁵⁰⁾ Over a follow-up out to 7 years, sST2 levels were predictive of total mortality (HR = 2.01; 95% CI, 1.15-3.51; $P=.01$). Soluble ST2 levels were also associated with risk of acute cellular rejection, with a significant difference between the top and bottom quartiles of sST2 levels in the risk of rejection ($P=.003$).

Pascual-Figal et al (2011), reported on 26 patients with post-cardiac transplantation with and an acute rejection episode.⁽⁵¹⁾ Soluble ST2 levels were measured during the acute rejection episode and compared with levels measured when acute rejection was not present. Soluble ST2 levels were higher during the acute rejection episode (130 ng/mL) than during the nonrejection period (50 ng/mL; $P=.002$). Elevated sST2 levels greater than 68 ng/mL had a positive predictive value of 53% and a negative predictive value of 83% for the presence of acute cellular rejection. The addition of sST2 levels to serum BNP resulted in incremental improvement in identifying rejection episodes.

Table 5. Summary of Key Nonrandomized Clinical Validity Study Characteristics

Study	Study Type	Country	Dates	Participants	Treatment	Follow-Up
Januzzi (2013) ⁵⁰	Retrospective	United States	NR	Post-cardiac transplantation	sST2 levels assessment (n=241)	Median 7.1 years
Pascual-Figal (2011) ⁵¹	Retrospective	Spain	2002-2007	Post-cardiac transplantation with acute rejection	sST2 levels assessment (n=26)	Median 3 months

NR: not reported, sST2: soluble suppression of tumorigenicity-2.

Table 6. Summary of Key Nonrandomized Clinical Validity Study Results

Study	Total Mortality	ST2 Levels	Positive Predictive Value	Negative Predictive Value
Januzzi (2013) ⁵⁰			NR	NR
HR (95% CI)	2.02 (1.16-3.52)	≥ 30 ng/mL at 7-year follow-up	NA	NA
<i>P-value</i>	.01	NR	NA	NA
Pascual-Figal (2011) ⁵¹			53%	83%
Rejection Episode	NR	130 ng/mL (IQR 60-238 ng/mL)	NA	NA
Nonrejection Period	NR	50 ng/mL (IQR 28-80 ng/mL)	NA	NA
<i>P-value</i>	NR	.002	NA	NA

HR: hazard ratio; NA: not applicable; NR: not reported; IQR: interquartile range.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from RCTs.

No RCTs were identified using sST2 levels that directed patient management in heart transplantation patients, and which assessed patient outcomes.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

No inferences can be drawn about the clinical utility of sST2 levels for patients with heart transplantation.

Section Summary: Use of Soluble Suppression of Tumorigenicity-2 sST2 in Post-Heart Transplantation Patients

Few studies are available, and they are observational and retrospective. No prospective studies were identified that provide high-quality evidence on the ability of sST2 levels to predict transplant outcomes. One retrospective study (N = 241) found that sST2 levels were associated with acute cellular rejection and mortality; another study (N = 26) found that sST2 levels were higher during an acute rejection episode than before rejection.

MEASUREMENT OF VOLATILE ORGANIC COMPOUNDS FOR HEART TRANSPLANT

Clinical Context and Test Purpose

The purpose of the measurement of volatile organic compounds in individuals with a heart transplant is to assess heart allograft rejection in a noninvasive manner.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest is individuals with heart transplant.

Interventions

The test being considered measures volatile organic compounds to assess allograft rejection.

Comparators

The following test is currently being used to diagnose heart allograft rejection: routine endomyocardial biopsy.

Outcomes

The general outcomes of interest are overall survival (OS), test validity, morbid events, and hospitalizations. Follow-up over months to years is necessary to monitor for signs of allograft rejection.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Study Selection Criteria

For the evaluation of the clinical validity of measuring volatile organic compounds, studies that met the following eligibility criteria were considered:

- Reported on the accuracy of the marketed version of the technology (including any algorithms used to calculate scores)
- Included a suitable reference standard (describe the reference standard)
- Individual/sample clinical characteristics were described
- Individual/sample selection criteria were described.

Observational Studies

The U.S. Food and Drug Administration approval of the Heartsbreath test was based on the results of the Heart Allograft Rejection: Detection with Breath Alkanes in Low Levels (HARDBALL) study sponsored by the National Heart, Lung, and Blood Institute.(52) The HARDBALL study was a 3-year, multicenter study of 1061 breath samples in 539 heart transplant patients. Before scheduled endomyocardial biopsy, patient breath was analyzed by gas chromatography and mass spectroscopy for volatile organic compounds. The amount of C4 to C20 alkanes and monomethylalkanes was used to derive the marker for rejection, known as the breath methylated alkane contour. The breath methylated alkane contour results were compared with subsequent biopsy results, as interpreted by two readers using the International Society for Heart and Lung Transplantation (ISHLT) biopsy grading system as the criterion standard for rejection.(22)

The authors of the HARDBALL study reported that the abundance of breath markers that measured oxidative stress was significantly greater in grade 0, 1, or 2 rejection than in healthy normal persons. In contrast, in grade 3 rejection, the abundance of breath markers of oxidative stress were found to be reduced—most likely due to accelerated catabolism of alkanes and methylalkanes that make up the breath methylated alkane contour. The authors also reported finding that in identifying grade 3 rejection, the negative predictive value (NPV) of the breath test (97.2%) was similar to endomyocardial biopsy (96.7%) and that the breath test could potentially reduce the total number of biopsies performed to assess for rejection in patients at low risk for grade 3 rejection. The sensitivity of the breath test was 78.6% vs 42.4% with biopsy. However, the breath test had lower specificity (62.4%) and a lower positive predictive value (PPV; 5.6%) in assessing grade 3 rejection than biopsy (specificity, 97%; PPV=45.2%). In addition, the breath test was not evaluated in grade 4 rejection.

Findings from the HARDBALL study were published by Phillips et al (2004). No subsequent studies evaluating the use of the Heartsbreath test to assess for graft rejection were identified in literature updates.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials.

No randomized controlled trials (RCTs) assessing the measurement of volatile organic compounds to diagnose cardiac allograft rejection were identified.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Because the clinical validity of measuring volatile organic compounds to assess cardiac allograft rejection has not been established, a chain of evidence support clinical utility cannot be constructed.

Section Summary: Measurement of Volatile Organic Compounds for Heart Transplant

A published study found that for identifying grade 3 (now grade 2R) rejection, the NPV of the breath test the study evaluated (97.2%) was similar to endomyocardial biopsy (96.7%), and the sensitivity of the breath test (78.6%) was better than that for biopsy (42.4%). However, the breath test had a lower specificity (62.4%) and a lower PPV (5.6%) in assessing grade 3 rejection than a biopsy (specificity, 97%; PPV=45.2%). The breath test was also not evaluated for grade 4 rejection. At present, no studies evaluating the clinical utility for the measurement of volatile organic compound testing for heart transplant have been identified.

DONOR-DERIVED CELL-FREE DNA TESTING FOR HEART TRANSPLANT

Clinical Context and Test Purpose

The purpose of donor-derived cell-free DNA (dd-cfDNA) testing in individuals with a heart transplant is to assess for allograft rejection.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest are individuals with heart transplants.

Interventions

The test being considered is dd-cfDNA testing to assess for allograft rejection (i.e., AlloSure, Prospera, myTAIHEART).

The AlloSure and Prospera tests report the fraction of dd-cfDNA, with both tests using a proposed high-risk of active transplant rejection cutoff of $\geq 0.15\%$. Clinical interpretation of alternate thresholds for quiescence ($< 0.12\%$), injury (0.20%) and severe injury (0.35%) have also been proposed.

The myTAIHEART test uses proprietary software to quantitatively genotype cfDNA in the patient's plasma after cardiac transplant, and distinguish dd-cfDNA originating from the engrafted heart from cfDNA originating from the recipient's native cells. Production of the myTAIHEART test was halted in 2020.

Comparators

The following test is currently being used to diagnose cardiac allograft rejection: routine endomyocardial biopsy.

Outcomes

The general outcomes of interest are OS, test validity, morbid events, and hospitalizations. Follow-up over months to years is needed to monitor for signs of allograft rejection.

Beneficial outcomes resulting from a true-negative test result are avoiding unnecessary subsequent biopsy. Harmful outcomes resulting from a false-positive result may include unnecessary biopsy or unnecessary treatment. Harmful outcomes from a false-negative result are increased risk of adverse transplant outcomes.

In a triage scenario, the test would need to identify precisely a group of individuals that could safely forgo biopsy; therefore, the sensitivity, NPV, and negative likelihood ratio are key test performance characteristics.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Study Selection Criteria

For the evaluation of the clinical validity of dd-cfDNA testing, studies that met the following eligibility criteria were considered:

- Reported on the accuracy of the marketed version of the technology (including any algorithms used to calculate scores)
- Included a suitable reference standard (describe the reference standard)
- Individual/sample clinical characteristics were described
- Individual/sample selection criteria were described.

AlloSure

Khush et al (2019) published performance characteristics for the AlloSure Heart dd-cfDNA test as assessed in the D-OAR prospective, multicenter registry study.(53) Patients already undergoing AlloMap testing for surveillance were eligible for inclusion; however following a protocol amendment, dd-cfDNA specimens were only obtained in patients with clinical suspicion of rejection and a planned for-cause biopsy after 2016 through 2018. The majority of dd-cfDNA samples (81%) were drawn in the first-year post-transplant. The D-OAR cohort included 841 biopsy-paired dd-cfDNA results, of which 587 were performed for routine surveillance of rejection. Overall, cell-mediated rejection and antibody-mediated rejection were biopsy-confirmed in 17 and 18 cases, respectively. The AUC for detecting acute rejection was 0.64 (95% CI, 0.52 to 0.75). At a 0.2% cutoff for dd-cfDNA, the sensitivity, specificity, PPV, and NPV for detection of acute rejection was 80%, 44%, 8.9%, and 97.1% respectively. For the subgroup of patients undergoing surveillance, the sensitivity, specificity, PPV, and NPV were 38.1%, 84.0%, 8.1%, and 97.3%, with a corresponding AUC of 0.61 (95% CI, 0.46 to 0.74). Among for-cause samples, the sensitivity, specificity, PPV, and NPV were 53.8%, 76.1%, 11.6%, and 96.6%, respectively. The study is limited by the protocol changes designed to increase the number of observed rejection events overall and low availability of concurrent dd-cfDNA results with respect to biopsy specimens (58%).

Richmond et al (2023) published data on pediatric (n=60) and adult (n=61) heart transplant recipients (median age, 24.3) prospectively enrolled at 8 participating centers from August 2016 to October 2017 and followed for up to 12 months.(82) All patients had samples from one or more endomyocardial biopsies post-transplantation with Allosure dd-cfDNA testing within 24

hours prior to biopsy. dd-cfDNA level was blinded to participants and investigators over the study period. Median dd-cfDNA was found to be significantly higher in the patients who had biopsy-defined allograft rejection (ACR or AMR) compared with healthy allograft participants (0.21% versus 0.09%, $p < .0001$). An area under the curve (AUC) analysis yielded an AUC of 0.78 using a pre-defined dd-cfDNA threshold of 14% and resulted in a test sensitivity of 67% and a specificity of 79% (NPV = 94% and PPV = 34%), a sub-group analysis satisfying patients into adult or pediatric patients found similar results (AUC of the adult cohort = 0.81; AUC of the pediatric cohort = 0.79).

Prospera Heart

Kim et al (2022) (54) assessed the clinical validity of the Prospera Heart dd-cfDNA test versus endocardial biopsy for prediction of acute heart transplant rejection. The study included 811 samples (703 prospectively collected and 108 retrospectively collected) from 223 heart transplant patients with a planned biopsy from 2 U.S. centers. The median patient age was 54 years and 27% were female. Race/ethnicity of the study population was: 54% White, 21% Hispanic, 12% Black, 6% Asian and 5% other race/ethnicity. The majority (91% [737/811]) of reference standard biopsies were conducted for surveillance, and median dd-cfDNA was lower in the surveillance samples (0.04%) than the for-cause samples (0.22%). The time from transplant to biopsy was 10 weeks, and the total prevalence of acute rejection was 9.0%. Median dd-cfDNA % was 0.58% in patients with acute rejection, although fractions varied according to rejection type/grade and were higher in those with antibody mediated rejection (median range 0.44% to 3.43%) than those with acute cellular rejection (median range 0.045% to 0.13%). In patients without acute rejection, dd-cfDNA % was 0.04. Diagnostic accuracy for 3 dd-cfDNA fractions were explored: 0.12%, 0.15% and 0.20%. At a cut-off of 0.12%, sensitivity was 86.6%, specificity was 72.0%, PPV was 23.4%, and NPV 98.2%. Corresponding values at a dd-cfDNA cut-off of 0.15% were 78.6%, 76.9%, 25.1% and 97.3%, and 78.6%, 82.1%, 30.3% and 97.5% at a dd-cfDNA cut-off of 0.20%. This resulted in an AUC for detection of acute rejection of 0.86 (95% CI 0.77 to 0.96). The optimal dd-cfDNA fraction for detection of heart transplant rejection has yet to be established. Limitations of the study include potential selection bias, as only patients with a scheduled biopsy were included in the study, and study authors noted that the prevalence of acute rejection in the study cohort was higher than in other cohorts.

A retrospective cohort study conducted by Rodgers et al (2023) compared dd-cfDNA testing with Allosure, which examines 405 single nucleotide polymorphisms (SNPs), to Prospera, which evaluates 13,292 SNPs, in 112 heart transplant patients. (83) Participants were enrolled from October 2020 to January 2022 and had a median age of 60 years (IQR, 47 to 65 years). Both tests used a dd-cfDNA threshold value of 15%. Testing with Allosure resulted in a low sensitivity (39%) and high specificity (82%) for identification of acute rejection; the Prospera test had similar characteristics with sensitivity at an identical 39% and a negligible difference in specificity (84%). Between-group comparisons showed no difference between the two tests in this small cohort. PPV with the Allosure test was 6.2% compared to 7% in Prospera testing ($p = .7$) and NPV was 98% for both tests ($p = .76$).

myTAIHEART

In a study funded by TAI Diagnostics, Inc., North et al (2020) performed a blinded clinical validation study on 158 matched pairs of endomyocardial biopsy-plasma samples collected from 76 volunteer adult and pediatric heart transplant recipients (ages 2 months or older, and 8 days or more post-transplant) between June of 2010 and Aug 2016 from 2 Milwaukee

transplant centers.(25) Based on acute cellular rejection grade as defined by the 2004 International Society for Heart and Lung Transplantation (ISHLT) classification, ROC analysis was performed to evaluate diagnostic accuracy across all possible dd-cfDNA % cutoffs. To maximize diagnostic accuracy, Youden's Index was used to select the optimal cutoff, found to correspond to a donor fraction value of 0.32%. Using this cutoff, clinical performance characteristics of the assay included a NPV of 100.00% for grade 2R or higher acute cellular rejection, with 100.00% sensitivity and 75.48% specificity; AUC for this analysis was 0.842, indicative of robust ability of the donor fraction assay to rule out 2R or greater acute cellular rejection for donor fraction values less than 0.32%. There was no statistically significant correlation of donor fraction with age. Donor fraction elevation can also be caused by other forms of injury to the donor heart such as acute cellular rejection 1R, acute antibody-mediated rejection (AMR), and presence of coronary artery vasculopathy (CAV), thereby requiring correlation of myTAIHEART results with other clinical indicators.

In study funded by a grant from the National Institutes of Health and TAI Diagnostics, Inc., Richmond et al (2019) assessed 174 post-cardiac transplant patients from 7 centers (ages 2.4 months-73.4 years) with myTAIHEART testing (before transplant; 1, 4, and 7 days following transplant; and at discharge from transplant hospitalization) using blinded analysis of biopsy-paired samples.(55) All the patients were followed for at least 1 year. dd-cfDNA was higher in acute cellular rejection 1R/2R (n = 15) than acute cellular rejection 0R (healthy) (n = 42) (p = .02); an optimal donor fraction threshold (0.3%) was determined by the use of ROC analysis, revealing an AUC of 0.81 with a sensitivity of 0.65, specificity of 0.93, and an NPV of 81.8% for the absence of any allograft rejection.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from RCTs.

No RCTs assessing the measurement of dd-cfDNA to diagnose cardiac allograft rejection were identified.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Because the clinical validity of measuring dd-cfDNA to assess for cardiac allograft rejection has not been established, a chain of evidence to support clinical utility cannot be constructed.

Section Summary: Measurement of Donor-Derived Cell-Free DNA for Heart Transplant

Studies measuring dd-cfDNA suggest that the dd-cfDNA fraction is elevated in acute rejection, but optimal fraction cut-offs for detection of acute rejection have not been established. Using dd-cfDNA thresholds ranging from 0.12% to 0.32% resulted in NPVs ranging from 82% to 98% and AUCs ranging from 0.61 to 0.86 in 3 studies. At present, no studies evaluating the clinical utility for the measurement of dd-cfDNA for heart transplant rejection have been identified.

GENE EXPRESSION PROFILING AND DONOR-DERIVED CELL-FREE DNA TESTING FOR HEART TRANSPLANT

Clinical Context and Test Purpose

The purpose of gene expression profiling (GEP) and donor-derived cell-free DNA (dd-cfDNA) testing in individuals with a heart transplant is to assess allograft rejection.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest are individuals with heart transplants.

Interventions

The test being considered is GEP to assess allograft rejection (i.e., AlloMap), used alone or in combination with AlloSure Heart dd-cfDNA testing. The combination of these tests is commercially marketed as HeartCare (CareDx).

AlloMap test results are reported on a scale from 0 to 40, with a proposed high-risk cutoff of ≥ 30 for patients < 6 months post-transplant and ≥ 34 for patients ≥ 6 months post-transplant. The HeartCare report provides the AlloMap score, AlloMap score variability, and AlloSure percent dd-cfDNA. Direct guidance for the combined interpretation of results is not provided in the HeartCare report, but potential clinical implications of concordant and discordant test scenarios have been proposed.(56)

Comparators

The following test is currently being used to diagnose cardiac allograft rejection: routine endomyocardial biopsy.

Outcomes

The general outcomes of interest are overall survival, test validity, morbid events, hospitalizations. Follow-up over months to years to monitor for signs of allograft rejection.

Beneficial outcomes resulting from a true-negative test result are avoiding unnecessary subsequent biopsy. Harmful outcomes resulting from a false-positive result may include unnecessary biopsy or unnecessary treatment. Harmful outcomes from a false-negative result are increased risk of adverse transplant outcomes.

In a triage scenario, the test would need to identify precisely a group of patients that could safely forgo biopsy; therefore, the sensitivity, NPV, and negative likelihood ratio are key test performance characteristics.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Study Selection Criteria

For the evaluation of clinical validity of GEP testing, studies were sought that met the criteria described in the first indication above.

AlloMap

Systematic Reviews

A TEC Assessment (2011) reviewed the evidence on the use of GEP using the AlloMap test.(57) The Assessment concluded that the evidence was insufficient to permit conclusions about the effect of the AlloMap test on health outcomes. Key evidence in the TEC Assessment is described below.

Nonrandomized Studies

Patterns of gene expression for development of the AlloMap test were studied in the Cardiac Allograft Rejection Gene Expression Observation (CARGO) study, which included 8 U.S. cardiac transplant centers enrolling 629 cardiac transplant recipients.(58) The study included discovery and validation phases. In the discovery phase, patient blood samples were obtained during the endomyocardial biopsy, and the expression levels of more than 7000 genes involved in immune responses were assayed and compared with the biopsy results. A subset of 252 candidate genes was identified from which, a panel of 11 genes was selected for evaluation. A proprietary algorithm is applied to the results of the analysis, producing a single score that considers the contribution of each gene in the panel.

The validation phase of the CARGO study, published by Deng et al (2006), was prospective, blinded, and enrolled 270 patients.(58) Primary validation was conducted using samples from 63 patients independent from discovery phases of the study and enriched for biopsy-proven evidence of rejection. A prospectively defined test cutoff value of 20 resulted in a sensitivity of 84% of patients with moderate/severe rejection but a specificity of 38%. Of note, in the “training set” used in the study, these rates were 80% and 59%, respectively. The authors evaluated the 11-gene expression profile on 281 samples collected at 1 year or more from 166 patients who were representative of the expected distribution of rejection in the target population (and not involved in discovery or validation phases of the study). When a test cutoff of 30 was used, the NPV (no moderate/severe rejection) was 99.6%; however, only 3.2% of specimens had grade 3 or higher rejection. In this population, grade 1B scores were found to be significantly higher than grade 0, 1A, and 2 scores but were similar to grade 3 scores.

A second prospective multicenter study, evaluating the clinical validity of GEP with the AlloMap test (CARGO II), was published by Crespo-Leiro et al (2016).(59) The study enrolled 499 heart transplant recipients who were undergoing surveillance for allograft rejection. The reference standard for rejection status was histologic grade from an endomyocardial biopsy performed on the same day as blood samples were collected. Blood samples needed to be collected 55 days or more post-transplant, more than 30 days after blood transfusion, more than 21 days after administration of prednisone 20 mg/day or more, and more than 60 days after treatment for a prior rejection. Patients had a total of 1579 eligible blood samples for which paired GEP scores, and endomyocardial biopsy rejection grades were available.

As in the original CARGO study, the proportion of cases of rejection was small. The prevalence of moderate-to-severe rejection (grade 2R/>3A) reported by local pathologists was 3.2%, which was reduced to 2.0% when confirmation from 1 or more other independent pathologist was required. At a GEP cutoff of 34, for patients who were at least 2 to 6 months posttransplant, the sensitivity of GEP for detecting grade 2R/>3A was 25.0% and the specificity was 88.7%. The PPV and NPV were 4.0% and 98.4%, respectively. Using the same cutoff of 34, for patients more than 6 months posttransplant, the sensitivity of GEP was 25.0% the specificity was 88.8%, the PPV was 4.3%, and the NPV was 98.3% The number of true-positives used in the above calculations was 5 (9.1%) of 55 for patients at least 2 to 6 months posttransplant and 6 (10.2%) of 59 for patients more than 6 months posttransplant.

Kanwar et al (2021) published data from the Outcomes AlloMap Registry (OAR) indicating that asymptomatic or active cytomegalovirus infection is associated with significantly higher AlloMap scores among heart transplant recipients compared to those without infection, even in the absence of acute rejection, potentially resulting in unnecessary biopsies among surveillance patients.(60) Donor-derived cell-free DNA levels measured by the AlloSure Heart test available for a small subset of samples (5.3%) were not significantly different between groups. The authors conclude that further assessment of the combined use of AlloMap and AlloSure scores is required to determine if this will improve differentiating infection-related from rejection-related immune activation. The combined use of these tests, commercially available as HeartCare (CareDx), is addressed in the following section.

HeartCare

The commercially available HeartCare (CareDx) test combines AlloMap GEP testing with AlloSure Heart measurement of percent dd-cfDNA. The combined use of GEP and dd-cfDNA testing for surveillance of acute rejection was assessed in a single-center, retrospective study conducted by Gondi et al (2021) between February 2019 and March 2020.(61) Patients (N=153) were required to be ≥ 55 days post-transplant, hemodynamically stable, ≥ 15 years of age, and single-organ recipients. The majority of patients were male (74.5%) and white (78.4%) with an average age of 54.5 years. Patients were assessed once monthly between 2 and 12 months, every 3 months between 12 and 24 months, and every 6 months between 24- and 36-months post-transplant. Pre-specified thresholds for GEP scores were ≥ 30 for patients < 6 months post-transplant and ≥ 34 for patients ≥ 6 months post-transplant. The pre-specified threshold for percent dd-cfDNA was $\geq 0.20\%$ based on a prior study of the AlloSure test by Khush et al (2019),(53) described in the following section. In patients < 6 months post-transplant, endomyocardial biopsy was performed regardless of test results. For patients ≥ 6 months post-transplant who received both GEP and dd-cfDNA testing, endomyocardial biopsy was canceled in patients with dd-cfDNA $< 0.20\%$ regardless of AlloMap score. In patients with positive AlloMap scores but negative dd-cfDNA, endomyocardial biopsy could be performed or deferred in favor of repeat dd-cfDNA testing. Among 495 samples, overall test result distributions were 59.6% for patients negative on both tests, 12.3% for patients positive by dd-cfDNA only, 22.6% for patients positive by GEP only, and 5.5% positive by both GEP and dd-cfDNA. The combined testing approach resulted in a 12.7% reduction (48 biopsies) in endomyocardial biopsy volume compared to GEP testing alone. Among the 172 biopsies performed, 2 patients with cell-mediated rejection were identified, with corresponding dual-positive tests. Two patients with antibody-mediated rejection were identified, with corresponding tests that were only positive by dd-cfDNA. The study is limited by its retrospective design, incomplete evaluation of performance characteristics, and lack of reporting on health outcomes.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials.

AlloMap

Randomized Controlled Trials

Kobashigawa et al (2015) published results of a pilot RCT evaluating the use of the AlloMap test in patients who were 55 days to 6 months post-transplant.(62) The trial design was similar to that of the Invasive Monitoring Attenuation through Gene Expression (IMAGE) RCT: 60 subjects were randomized to rejection monitoring with AlloMap or with endomyocardial biopsy at prespecified intervals of 55 days and 3, 4, 5, 6, 8, 10, and 12 months posttransplant. The threshold for a positive AlloMap test was set at 30 for patients 2 to 6 months posttransplant and 34 for patients after 6 months posttransplant, based on data from the CARGO study. Endomyocardial biopsy outside of the scheduled visits was obtained in either group if there was clinical or echocardiographic evidence of graft dysfunction and for the AlloMap group if the score was above the specified threshold. The incidence of the primary outcome at 18 months posttransplant (composite outcome of first occurrence of any of the following: death or retransplant, rejection with hemodynamic compromise, or allograft dysfunction due to other causes) did not differ significantly between the AlloMap and biopsy groups (10% vs 17%; $p=.44$). The number of biopsy-proven rejection episodes (International Society for Heart and Lung Transplantation grading system $\geq 2R$) within the first 18 months did not differ significantly between groups (3 in the AlloMap group vs 1 in the biopsy group; $p=.31$). Of the rejections in the AlloMap group, one was detected after an elevated routine AlloMap test, while two were detected after patients presented with hemodynamic compromise. As in the IMAGE study described above, a high proportion of rejection episodes were detected by clinical signs or symptoms (however, this study had only three rejection episodes in the AlloMap group).

In 2010, the results of the IMAGE study were published.(63,64) This was an industry-sponsored, nonblinded, noninferiority randomized controlled trial (RCT) that compared outcomes in 602 patients managed with the AlloMap test ($n=297$) or with routine endomyocardial biopsies ($n=305$). The study included adults from 13 centers who underwent cardiac transplantation between 1 and 5 years prior to participating in the study, were clinically stable, and had a left ventricular ejection fraction of at least 45%. To increase enrollment, the study protocol was later amended to include patients who had undergone transplantation between 6 months and 1 year prior to participating in the study; this subgroup ultimately comprised only 15% of the final sample ($n=87$). Each transplant center used its own protocol for determining the intervals for routine testing. At all sites, patients in both groups underwent clinical and echocardiographic assessments in addition to the assigned surveillance strategy. According to the study protocol, patients underwent biopsy if they had signs or symptoms of rejection or allograft dysfunction at clinic visits (or between visits) or if the echocardiogram showed a left ventricular ejection fraction decrease of at least 25% compared with the initial

visit. Additionally, patients in the AlloMap group underwent biopsy if their test score was above a specified threshold; however, if they had two elevated scores with no evidence of rejection found on two previous biopsies, no additional biopsies were required. The AlloMap test score varied from 0 to 40, with higher scores indicating a higher risk of transplant rejection. The investigators initially used 30 as the cutoff for a positive score; the protocol was later amended to use a cutoff of 34 to minimize the number of biopsies needed. Fifteen patients in the AlloMap group and 26 in the biopsy group did not complete the trial.

The primary outcome was a composite variable: (1) the first occurrence of rejection with hemodynamic compromise; (2) graft dysfunction due to other causes; (3) death; or (4) retransplantation. Use of the AlloMap test was considered noninferior to the biopsy strategy if the 1-sided upper boundary of the 95% confidence interval (CI) for the hazard ratio (HR) comparing the 2 strategies was less than the prespecified margin of 2.054. The margin was derived using the estimate of a 5% event rate per year in the biopsy group, taken from published observational studies, and allowing for an event rate of up to 10% per year in the AlloMap group.

According to Kaplan-Meier analysis, the 2-year event rate was 14.5% in the AlloMap group and 15.3% in the biopsy group. The corresponding HR was 1.04 (95% CI, 0.67 to 1.68). The upper boundary of the CI of the HR (1.68) fell within the prespecified noninferiority margin (2.054); thus, GEP was considered noninferior to endomyocardial biopsy. Death from all causes, a secondary outcome, did not differ significantly between groups. There were 13 (6.3%) deaths in the AlloMap group and 12 (5.5%) in the biopsy group ($p=.82$). During follow-up, there were 34 treated episodes of graft rejection in the AlloMap group. Only 6 (18%) of the 34 patients with graft rejection presented solely with elevated AlloMap scores. Twenty (59%) patients presented with clinical signs/symptoms and/or graft dysfunction on echocardiogram, and 7 patients had an elevated AlloMap score plus clinical signs/symptoms with or without graft dysfunction on echocardiogram. In the biopsy group, 22 patients were detected solely due to an abnormal biopsy.

A total of 409 biopsies were performed in the AlloMap group and 1249 in the biopsy group. Most of the biopsies in the AlloMap group (67%) were performed because of elevated gene profiling scores. Another 17% were performed due to clinical or echocardiographic manifestations of graft dysfunction, and 13% were performed as part of routine follow-up after treatment for rejection. There was 1 (0.3%) adverse event associated with biopsy in the AlloMap group and 4 (1.4%) in the biopsy group. In terms of quality of life, the physical health and mental health summary scores of the 12-Item Short Form Health Survey (SF-12) were similar in the 2 groups at baseline and did not differ significantly between groups at 2 years.

A limitation of the trial was that the threshold for a positive AlloMap test was changed partway through the study; thus, the optimal test cutoff remains unclear. Moreover, the trial was not blinded, which could have affected treatment decisions based on clinical findings, such as whether to recommend a biopsy. In addition, the study did not include a group that only received clinical and echocardiographic assessment, so the value of AlloMap testing beyond that of clinical management alone cannot be determined. The uncertain incremental benefit of the AlloMap test is highlighted by the finding that only 6 of the 34 treated episodes of graft rejection detected during follow-up in the AlloMap group were initially identified solely due to an elevated GEP score. Since 22 episodes of asymptomatic rejection were detected in the biopsy group, the AlloMap test does not appear to be a sensitive test, possibly missing more than half

of the episodes of asymptomatic rejection. Because clinical outcomes were similar in the 2 groups, there are at least 2 possible explanations: the clinical outcome of the study may not be sensitive to missed episodes of rejection, or it is not necessary to treat asymptomatic rejection. In addition, the trial was only statistically powered to rule out more than a doubling of the rate of the clinical outcome, which some may believe is an insufficient margin of noninferiority. Finally, only 15% of the final study sample had undergone transplantation less than 1 year before study participation; therefore, findings might not be generalizable to the population of patients 6 to 12 months post-transplant.

HeartCare

Direct evidence of clinical utility was not identified for the HeartCare test.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Because the clinical validity of GEP testing, alone or in combination with dd-cfDNA testing, to assess for cardiac allograft rejection has not been established, a chain of evidence to support clinical utility cannot be constructed.

Section Summary: Gene Expression Profiling for Heart Transplant

AlloMap

The 2 studies (CARGO, CARGO II) examining the diagnostic performance of GEP using the AlloMap test for detecting moderate or severe rejection were flawed by lack of a consistent threshold (i.e., 20, 30, or 34) for determining positivity and by a small number of positive cases. In the available studies, although the NPVs were relatively high (i.e., at least 88%), the performance characteristics were calculated based on the detection of 10 or fewer cases of rejection each. Moreover, the PPV in the CARGO II study was only 4.0% for patients who were at least 2 to 6 months post-transplant and 4.3% for patients more than 6 months post-transplant. The ability of the AlloMap test to differentiate between infection-related and rejection-related graft injury has also been called into question.

The most direct evidence on the clinical utility of GEP using the AlloMap test comes from a large RCT comparing a GEP-directed strategy with an endomyocardial biopsy-directed strategy for detecting rejection; it found that the GEP-directed strategy was noninferior. However, given the high proportion of rejection episodes in the GEP-directed strategy group detected by clinical signs/symptoms, the evidence is insufficient to determine that health outcomes are improved because of the uncertain incremental benefit of GEP. In addition, a minority of subjects assessed were in the first year post transplant. Results from a pilot RCT would suggest that GEP may have a role in evaluating for heart transplant rejection beginning at 55 days post-transplant, but the trial was insufficiently powered to permit firm conclusions about the noninferiority of early GEP use.

Subsection Summary: Gene Expression Profiling with Donor-Derived Cell-Free DNA Testing for Heart Transplant

HeartCare

One retrospective study assessing the combined use of GEP testing with AlloMap and dd-cfDNA testing with AlloSure Heart reported a 12.7% reduction in endomyocardial biopsy volume when combined testing was used compared to AlloMap alone. However, this observation is limited by a lack of reporting on long-term health outcomes and incomplete diagnostic performance assessment for combined testing.

DONOR-DERIVED CELL-FREE DNA (ALLOSURE) IN CONJUNCTION WITH GENE EXPRESSION PROFILING (ALLOMAP) TESTING FOR HEART TRANSPLANT

According to CareDx, HeartCare provides a comprehensive assessment of graft rejection by combining AlloMap Heart gene expression profiling with Allosure Heart dd-cfDNA. AlloMap Heart assesses immune quiescence, while AlloSure Heart detects graft injury.(65) The laboratory claims that when AlloMap and AlloSure are used in combination (HeartCare assay) the panel is better at detecting heart transplant rejection than when they are used as individual tests. AlloSure detects graft injury and rejection by measuring the amount of dd-cfDNA in the blood. When graft injury occurs, dd-cfDNA levels rise. The assay determines the level of dd-cfDNA using single nucleotide polymorphisms (SNPs) to distinguish cell-free DNA (cfDNA) from the allograft and the transplant recipient. A high dd-cfDNA fraction (associated with an increased risk of rejection) was defined as $\geq 0.2\%$ dd-cfDNA. Key intervals in the use of the HeartCare assay are recommended in order to track changes in the levels overtime. The laboratory provides a Heart Allograft Routine Testing Schedule (HARTS), which recommends monthly testing for the first year, quarterly for year 2 and 3, and biannually thereafter. Based on a review of the laboratory website, it was not clear how each test result for this comprehensive solution are provided: whether provided separately, together, or integrated.

Kanwar et al (2020) discussed cytomegalovirus (CMV) infection implications in the pathogenesis of allograft rejection, which is reflected by high gene expression profiling (GEP) scores.(66) The effect of CMV infection on serial GEP scores in the absence of acute cellular rejection (ACR) is unknown. Data from 14,985 samples collected from 2288 heart transplant (HT) recipients enrolled in Outcomes AlloMap® Registry (OAR) were analyzed. The empirical distribution of GEP scores at the visit closest to CMV infection were compared to GEP scores obtained at other visits via Kolmogorov-Smirnov tests. AlloSure measurements from those who underwent concurrent dd-cfDNA testing were also reviewed. Three hundred and eleven samples were collected from 213 patients with CMV infection at the time of GEP testing. The mean time to CMV infection from transplant was 346 days. Median GEP score at the time of diagnosis of CMV infection was significantly higher when compared to GEP score obtained at other visits (34 v 30, $p < 0.0001$) (Figure 1). For patients with samples assessed via AlloMap and AlloSure, there was no significant difference in the distribution of AlloSure measurements at the time of CMV infection. CMV infections in HT recipients are associated with an increase in GEP score independent of associated ACR. The likely mechanism is immune activation/modulation of one or more of the 11 genes in the GEP signature. Hence, CMV status should be considered when clinically interpreting high GEP scores. CMV infection in HT recipients is associated with an increase in AlloMap score, whereas AlloSure results do not appear to be impacted. This information should be considered when clinically interpreting

abnormal/high AlloMap scores in HT recipients. Disclosure indicates that authors are affiliated with CareDx.

Purewal et al (2020) performed a retrospective study of data from Allosure cfDNA and AlloMap GEP assays (CareDx).⁽⁶⁷⁾ Cell-free DNA results were collected from the DOAR Registry as well as cfDNA and GEP from clinical testing. Test sensitivity (SN), specificity (SP), positive predictive values (PPV) and negative predictive values (NPV) were able to be derived from evaluating the combination of these results and comparing them to endomyocardial biopsy (EMBx), which is the gold standard for acute cellular rejection (ACR) after heart transplantation (HT). A threshold of 0.2% for cfDNA was used. The utility of both non-invasive assays in combination were also assessed. One hundred sixty-eight patients (total 422 samples) with 67 patients from the initial D-OAR cohort were included. This cohort was predominantly male (72%) and Caucasian (60%) with a median (IQR) 316 (238 - 484) days from transplant to cfDNA test. A SN of 0.78, SP 0.75, PPV 0.30 and NPV of 0.96 to detect ACR was found. Among patients who had EMBx, GEP and cfDNA, 70% had concordant cfDNA and GEP results (Fig 1A). Of those with a positive cfDNA but negative GEP, none had significant (<2R) rejection. After both assays were available, there were 101 patients who had surveillance 237 times with both tests. In 85% (n=201) both tests were below threshold and, of these, 97% had no biopsy. When monitoring patients with both cfDNA and GEP, the most common result was that both were below threshold. When the results were discordant and a biopsy was performed, no treatable ACR was observed. Endomyocardial biopsy is the gold standard for acute cellular rejection after heart transplantation. With the commercial introduction of gene-expression profiling (GEP) and more recently, donor-derived cell-free DNA (cfDNA), the predictive ability of cfDNA and its utility in combination with GEP are not well known. Disclaimers indicated authors are affiliated with CareDx.

Although HeartCare is intended to assess the probability of allograft rejection in a particular patient, biopsy is still necessary to confirm and establish the type of active rejection in affected patients (68). It has been proposed that these tests be used for serial monitoring in order to detect new onset injury or rejection prior to clinical symptoms, however the optimal time interval has yet to be established (69). The clinical utility for the use of dd-cfDNA to evaluate transplant rejection has yet to be established. There is insufficient evidence to support the accuracy and clinical utility of donor-derived cell free DNA for assessing and monitoring the probability of allograft rejection in heart transplant patients.

Section Summary: Donor-Derived Cell-Free DNA (Allosure) in Conjunction with Gene Expression Profiling (Allomap) Testing for Heart Transplant

Published data comparing the combination of dd-cfDNA technology with gene expression profiling tests (HeartCare, CareDx) to current methods are lacking. Unbiased peer reviewed scientific studies are limited in the published literature. There were no studies identified to determine clinical utility or that the use of this technology in identifying cardiac transplant rejection leads to improved health outcomes. Further research is needed to determine the effect that the technology has on overall patient outcomes. Direct evidence for clinical utility is lacking.

DONOR-DERIVED CELL-FREE DNA TESTING FOR RENAL TRANSPLANT

Clinical Context and Test Purpose

The purpose of dd-cfDNA testing in individuals with renal transplant who are undergoing surveillance or have clinical suspicion of allograft rejection is to assess for allograft rejection.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest are individuals with renal transplants and clinical suspicion of allograft rejection.

Clinical suspicion of allograft rejection may be indicated by clinical symptoms (e.g., pain) or dynamic changes in laboratory parameters.

Allograft dysfunction is typically asymptomatic and has a broad differential, including graft rejection. Diagnosis and rapid treatment are recommended to preserve graft function and prevent loss of the transplanted organ.

Interventions

The test being considered is measurement of dd-cfDNA to assess renal allograft rejection (i.e., AlloSure or Prospera).

Various clinical pathways have been proposed for these tests. Use of the Prospera test is recommended when there is clinical suspicion of active rejection and for regular surveillance of subclinical rejection.(35) In a surveillance scenario, regular testing is recommended at 1, 2, 3, 4, 6, 9, and 12 months after renal transplant or most recent rejection. Thereafter, the test should be repeated quarterly. The proportion of dd-cfDNA relative to total cfDNA is reported, with detection of $\geq 1\%$ dd-cfDNA indicating increased risk for active rejection. The percent dd-cfDNA change between tests is also reported.(36) In the surveillance scenario, individuals with a negative result may avoid biopsy and it is recommended that a positive test result is incorporated with clinical findings to determine whether a biopsy is indicated. When there is clinical suspicion of rejection, testing is recommended as an adjunct to biopsy for treatment response monitoring, or as a rule-out test for biopsy.

For the AlloSure test, various dd-cfDNA thresholds are suggested depending on the clinical scenario and include the detection of antibody-mediated rejection (ABMR) in patients with donor-specific antibodies (DSA), the detection of "likely" active rejection, the prediction of adverse outcomes as an adjunct to biopsy-confirmed T cell-mediated (TCMR) 1A/borderline rejections, and for the exclusion of active rejection.(70) A routine testing schedule is also recommended, and details regarding its clinical rationale have been published.(71)

Comparators

The following test is currently being used to confirm clinical suspicion of allograft rejection: renal biopsy. The adoption of protocol (i.e., surveillance) biopsies varies across transplant centers and its use is not standardized.

Clinical suspicion of allograft rejection may be indicated by physical symptoms and/or dynamic changes in laboratory parameters (e.g., serum creatinine, estimated glomerular filtration rate [eGFR], DSA).

Outcomes

The general outcomes of interest are overall survival, test validity, morbid events, hospitalizations. Follow-up over months to years is needed to monitor for signs of allograft rejection.

For a primary kidney transplant, graft survival at 1 year is 94.7%; at 5 years, graft survival is 78.6%.(29)

Beneficial outcomes resulting from a true-negative test result are avoiding unnecessary subsequent biopsy. Harmful outcomes resulting from a false-positive result may include an unnecessary biopsy or unnecessary treatment. Harmful outcomes from a false-negative result are increased risk of adverse transplant outcomes.

In a triage scenario, the test would need to identify precisely a group of patients that could safely forgo biopsy; therefore, the sensitivity, NPV, and negative likelihood ratio are key test performance characteristics.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Study Selection Criteria

For the evaluation of clinical validity of dd-cfDNA testing, studies that met the criteria described in the first indication above.

Meta-Analyses

Two meta-analyses were identified which assessed the clinical validity of dd-cfDNA testing.(84,85) Both studies quantitatively synthesized the findings from 9 observational studies to determine the diagnostic accuracy of dd-cfDNA as a potential marker of graft rejection following kidney transplantation. Xiao et al (2021) calculated a pooled sensitivity of 0.70 (95% CI, 0.57-0.81; I2, 65) and specificity of 0.78 (0.70-0.84; I2, 75) from 6 studies evaluating the diagnostic accuracy of dd-cfDNA for any rejection episode.(85) The area under the receiver operating characteristics curve (AUC) was 0.81 (95% CI, 0.77 to 0.84; I2, 65) with an overall diagnostic odds ratio (DOR) of 8.18 (95% CI, 5.11 to 13.09). Similar pooled estimates were calculated for 5 studies discriminating AMR. The authors reported a pooled sensitivity of 0.84 (95% CI, 0.75 to 0.90; I2, 0) and a specificity of 0.80 (95% CI, 0.75 to 0.84; I2, 4) with an AUC of 0.89 (95% CI, 0.86 to 0.91) and overall DOR of 20.48 (95% CI, 10.76 to 38.99). Overall, the authors found greater value in dd-cfDNA as a biomarker for AMR in patients with suspected renal dysfunction than in discriminating a main rejection episode and cite the need for more large-scale, prospective research on the topic.

Wijvliet et al (2020) performed stratified analyses of dd-cfDNA fraction and calculated pooled median estimates in the following patient groups: patients, patients without rejection at indication biopsy, patients with pure T cell-mediated rejection, and patients with a component AMR.(84) In stable patients (n=1149; 5 studies), the median dd-cfDNA fraction was 0.29%

(95% CI, 0.21% to 0.45%) and in the AMR group (n=89; 6 studies) the average was nearly 10 times greater (2.5%; 95% CI, 1.4% to 2.9%). In T cell-mediated rejection patients (n=35; 4 studies), the weighted median was found to be 0.27% (95% CI, 0.26% to 2.69%) and in patients without rejection (n=225; 4 studies) the weighted median was 0.57% (95% CI, 0.30 to 0.67). The authors also calculated the weighted median differences in medians (WMDMs) between groups and found that median dd-cfDNA fractions were significantly higher in patients with AMBR than in patients without rejection (1.89%; 95% CI, 1.02 to 2.6), stable patients (2.3%; 95% CI, 1.8 to 2.69). However, no significant difference was observed for WMDMs between AMR patients and T cell-mediated rejection patients or for comparing T cell-mediated rejection to stable patients. This review had moderate heterogeneity for most between-group comparisons. Overall, higher dd-cfDNA fractions were found in patients with AMR than in individuals without rejection or stable patients, but a less clear relationship was established for T cell-mediated rejection and other investigated subgroups.

Observational Studies

AlloSure

Development of the AlloSure test was conducted in the multicenter prospective Circulating Donor-Derived Cell-Free DNA in Blood for Diagnosing Acute Rejection in Kidney Transplant Recipients (DART) study by Bloom et al (2017), which both recruited patients who were less than 3 months after renal transplant (n=245) and recruited renal transplant patients requiring a biopsy for suspicion of graft rejection (n=139).⁽⁶⁹⁾ For the primary analysis, active rejection was defined as the combined categories of T cell-mediated rejection, acute/active antibody-mediated rejection, and chronic/active antibody-mediated rejection as defined by the Banff working groups. Only patients undergoing biopsy were considered; further exclusion of biopsies which were not for cause, had an inadequate or incomplete collection of biopsies or corresponding blood samples, or had prior allograft in situ resulted in the main study cohort (N=102 patients, 107 biopsies). Within this population, acute rejection was noted in 27 patients (27 biopsies). After statistical analysis accounting for multiple biopsies from the same patient, the threshold dd-cfDNA fraction corresponding to acute rejection was set to 1.0% or higher. In the main study group, this resulted in a sensitivity of 59% (95% CI, 44% to 74%) and specificity of 85% (95% CI, 79% to 81%) for detecting active rejection vs no rejection. Using the original data set including all biopsies performed for clinical suspicion of rejection, 58 cases of acute rejection were diagnosed in 204 biopsies (170 patients). This PPV was 61% and the NPV 84%. Biopsies performed for surveillance (n=34 biopsies) were excluded from analysis in this study as only one biopsy for surveillance demonstrated acute rejection. Study limitations included the absence of a validation data set.

Huang et al (2019) conducted a smaller single center that recruited 63 renal transplant patients with suspicion of rejection that had AlloSure assessment of dd-cfDNA within 30 days of an allograft biopsy.⁽⁷²⁾ Median years from transplant to dd-cfDNA measurement was 2.0 (interquartile range, 0.3 to 6.5). Within this population, biopsy found acute rejection in 34 (54%) of patients; 10 (15.9%) were cell-mediated only, 22 (25.4%) were antibody-mediated only, and 2 (3.2%) were mixed cell-mediated and antibody-mediated. In contrast to the study by Bloom et al (2017), the optimal threshold for a positive dd-cfDNA result was identified as $\geq 0.74\%$. For the outcome of any rejection (i.e., cell-mediated, antibody-mediated, or mixed), use of this threshold was associated with an overall sensitivity of 79.4%, specificity of 72.4%, PPV of 77.1%, and NPV of 75.0%. Discrimination of rejection differed by biopsy findings, however. For the subgroup of patients with antibody-mediated rejection, the sensitivity was 100%, specificity

was 71.8%, PPV was 68.6%, and NPV was 100%. The dd-cfDNA test did not discriminate rejection in patients with cell-mediated rejection, as evidenced by an AUC of 0.43 (95% CI, 0.17 to 0.66). Major limitations of this study is its small sample size and single-center setting.

Stites et al (2020) assessed clinical outcomes in 79 patients diagnosed with TCMR 1A/borderline rejection with simultaneous AlloSure assessment of dd-cfDNA across 11 centers between June 2017 and May 2019.(73) Timing of testing with respect to the date of transplantation was not reported. Elevated levels of dd-cfDNA ($\geq 0.5\%$) were detected in 42 (53.2%) patients. No statistically significant differences between dd-cfDNA distributions when stratified by protocol versus for-cause biopsies was detected ($p = .7307$). Elevated levels of dd-cfDNA were associated with adverse clinical outcomes compared to patients with low levels ($< 0.5\%$), including decline in eGFR (8.5% versus 0%; $p = .004$), de novo DSA formation (40% versus 2.7%; $p < .0001$), and future or persistent rejection (21.4% versus 0%; $p = .003$). The authors hypothesize that the use of dd-cfDNA may complement histological evaluation and risk stratify patients with TCMR 1A or borderline rejection identified on biopsy and propose the use of reference ranges as opposed to absolute dd-cfDNA cutoff thresholds.

Additional analyses of the DART study have reported on associations between first-year AlloSure dd-cfDNA fraction or serial variability and subsequent eGFR decline (Sawinski et al [2021]),(74) and combined use of dd-cfDNA and DSA testing to diagnose active antibody-mediated rejection (Jordan et al [2018], Mayer et al [2021]).(69,75)

Puliyanda et al (2021) conducted a prospective study of 67 pediatric renal transplant recipients enrolled across 2 medical centers between 2017 and 2019.(76) Patients had a median age of 11 years (interquartile range [IQR], 4 to 13) and median time post-transplant to first AlloSure dd-cfDNA measurement was 55.6 months. Nineteen patients (28.4%) received dd-cfDNA testing in the absence of clinical suspicion of rejection. Median dd-cfDNA scores in the surveillance versus for-cause cohorts were 0.37% (IQR, 0.19% to 1.10%) and 0.47% (IQR, 0.24% to 2.15%), respectively. Among patients undergoing surveillance, 26.3% (5/19 patients) had a dd-cfDNA score $>1\%$ with biopsies indicating 4 cases of antibody-mediated rejection and 1 case of mixed rejection. Among patients with clinical suspicion of rejection, 43.8% (21/48 patients) had dd-cfDNA scores $>1\%$. All for-cause biopsies showed evidence of rejection, including 11 cases of antibody-mediated rejection, 2 cases of T cell-mediated rejection, and 8 cases of mixed rejection. An additional 7 patients with clinical suspicion of rejection underwent biopsy despite dd-cfDNA scores $< 1\%$, revealing 4 cases without rejection, 1 case with antibody-mediated rejection, 1 case with cell-mediated rejection, and 1 case of mixed rejection. Among all patients with biopsy-matched results (33/67), dd-cfDNA $>1\%$ was associated with a sensitivity of 86% and specificity of 100%, with a corresponding AUC of 0.996 ($p = .002$). No significant difference in serum creatinine change from baseline to testing was identified for those with rejection compared to those without. The study is limited by the small sample size and lack of biopsy-matched data for a complete assessment of false negatives. The authors also note that the 1% dd-cfDNA cutoff threshold was used based on prior studies in adults and it is unclear if this is appropriate for the pediatric population. Additionally, the authors suggest that relative increases in dd-cfDNA, as opposed to absolute values, may be more valuable in the pediatric population, given that appropriate cutoff thresholds may depend on child age and size.

Dandamudi et al (2022) conducted a prospective study of 57 pediatric renal transplant recipients enrolled in a single center from 2013 to 2019.(86) Patients had a median age of 14 years (IQR, 7.5 to 16) and time post-transplantation to first Allosure dd-cfDNA measurement was within 30 days and through 12 months post-transplantation. The authors attempted to correlate dd-cfDNA scores to biopsy-proved T cell-mediated rejection (including sub-clinical rejection). Twenty-two of the patients had biopsy-proven rejection, and cfDNA median levels were higher in these patients (0.91%, IQR, 0.54% to 1.2%) than in the patients without biopsy-proven rejection (median, 0.22%; IQR, 0.14% to 0.45%; $p < .001$). An area under the receiver operating characteristic curve value of .82 (95% CI, 0.71 to 0.93) was found between dd-cfDNA level and biopsy-proven rejection. Using a cut-off of 1%, cfDNA had a high specificity (96%; 95% CI, 90% to 99%) and low sensitivity (33%; 95% CI, 19% to 52%). A lower cut-off of 0.5% dd-cfDNA had a lower specificity (79%; 95% CI, 69% to 87%) but had a higher sensitivity (78%; 95% CI, 59% to 89%).

Bu et al (2022) evaluated data from 1092 kidney transplant recipients at 7 centers from June 2016 to January 2020 as part of the ADMIRAL study (NCT0456605).(87) All patients were monitored with Allosure dd-cfDNA as part of their standard care. A total of 1092 adult kidney transplant recipients (mean age 49.5 years) were followed for a period of up to 3 years to determine the association of dd-cfDNA with evidence of allograft rejection identified histologically. Using a cfDNA threshold of 0.5%, the authors found an increase in the risk of the development of donor-specific antibodies (hazard ratio [HR], 2.7). Having a dd-cfDNA result of more than 0.5% on more than 1 test predicted a reduction in eGFR over 3 years (HR, 1.97). The presence of allograft rejection was established using results from 203 patients who had a biopsy to pain with cfDNA results. Amongst patients with no rejection on biopsy, a median dd-cfDNA level of 0.23% (IQR, 0.19% to 0.64%) was lower than that observed in individuals with biopsy-defined cellular or humoral rejection (1.6%; IQR, 0.68% to 2.6%; $p < .0001$). Median dd-cfDNA levels had an area under the receiver operating characteristic curve (AUROC) of 0.8 for graft rejection (95% CI, 0.72 to 0.87) and was found to be more predictive than the AUROC of median creatinine levels in this sample of patients. Performance characteristics of the Allosure test at a dd-cfDNA threshold of 0.5% resulted in a sensitivity of 78% and specificity of 71%; using a dd-cfDNA cut-off of 1.0 reduced the test sensitivity to 58% but improved the specificity to 82%.

Huang and colleagues (2023) reported a retrospective cohort study of all kidney transplant patients at a single center who received testing with Allosure.(88) A total of 317 individuals who underwent kidney transplantation were included in this study (median age, 55 years) and were defined as either low ($< 0.5\%$, $n=239$), moderate (0.5% to $< 1\%$, $n=43$), or high ($\geq 1\%$, $n=35$) based on dd-cfDNA threshold levels. The rejection rate was established by comparing the 62 participants who underwent a biopsy; patients in the low dd-cfDNA group had a rejection rate of only 5% which was statistically less than that observed in the high dd-cfDNA group (17%; $p=.01$) but did not vary significantly in the moderate dd-cfDNA group (12%; $p=.13$). Although each group did not experience a significant change in eGFR from baseline levels, a linear mixed-effects model of eGFR over time found that dd-cfDNA category had a significant interaction when comparing both the moderate to low ($p=.005$) and low to high ($p=.048$) after adjustments for age, donor type, and history of donor-specific antibodies.

Prospera Kidney

Sigdel et al (2019) discussed a study which used a novel single nucleotide polymorphism (SNP)-based massively multiplexed PCR (mmPCR) methodology to measure dd-cfDNA in various types of renal transplant recipients for the detection of allograft rejection/injury without prior knowledge of donor genotypes.(77) A total of 300 plasma samples (217 biopsy-matched: 38 with active rejection (AR), 72 borderline rejection (BL), 82 with stable allografts (STA), and 25 with other injury (OI)) were collected from 193 unique renal transplant patients; dd- cfDNA was processed by mmPCR targeting 13,392 SNPs. Median dd-cfDNA was significantly higher in samples with biopsy-proven AR (2.3%) versus BL (0.6%), OI (0.7%), and STA (0.4%) ($p < 0.0001$ all comparisons). The SNP-based dd-cfDNA assay discriminated active from non-rejection status with an area under the curve (AUC) of 0.87, 88.7% sensitivity (95% CI, 77.7–99.8%) and 72.6% specificity (95% CI, 65.4–79.8%) at a prespecified cutoff (>1% dd-cfDNA). Although Prospera was not named anywhere in the article, a disclaimer indicated that many of the authors were affiliated with Natera, and the cell-free DNA was input into library preparation using the Natera Library prep kit. The authors concluded that this study validates the use of dd-cfDNA in the blood as an accurate marker of kidney injury/rejection across a range of pathologies with acute and chronic findings. However, several important limitations were noted in the study. Aside from the retrospective nature of the study in a single institution there were key differences in the groups. The acute rejection group contained mostly for-cause biopsies whereas the non-rejection group contained mostly surveillance biopsies. This is important because the incidence of rejection is known to be significantly greater in for-cause biopsies.(78) Further data are needed, ideally in a prospective real-world cohort, to confidently evaluate the performance of this test.

Sigdel et al (2019) also assessed the performance characteristics of eGFR, which was calculated as a function of serum creatinine with adjustments for age, sex, and race based on the Modification of Diet in Renal Disease (MDRD) Study equation.(77) At a cutoff threshold of < 60 , the sensitivity and specificity for eGFR were lower compared to dd-cfDNA, at 67.8% (95% CI, 51.3% to 84.2%) and 65.3% (95% CI, 57.6% and 73.0%), respectively, with a corresponding AUC of 0.74 (95% CI, 0.66 to 0.83). However, the relevance of absolute eGFR measurements is limited as dynamic changes in laboratory parameters (e.g., serum creatinine elevation, eGFR decline) are used to flag impaired kidney function in clinical practice in the transplant population. Separate eGFR estimates in the for-cause subgroup were not reported. Major limitations of this study include its retrospective design and single-center setting. While the dd-cfDNA cutoff was prespecified, it was based on prior studies of the AlloSure test and may not be optimized for Prospera.

Bunnapradist et al (2021) noted that while % dd-cfDNA is a promising noninvasive biomarker for detecting renal allograft rejection, levels can be artificially depressed by high levels of circulating cfDNA, which may be observed in patients who are obese, have recently undergone surgery, have medical complications, or receive certain medications, potentially leading to false-negative results.(79) The authors suggest that a combination of dd-cfDNA fraction and absolute quantity thresholds may improve the sensitivity of allograft rejection while maintaining high specificity.

Preliminary results from the ongoing Trifecta study (NCT04239703) published by Halloran et al (2022) provide assessment of combined dd-cfDNA fraction and absolute values for prediction of active kidney allograft rejection.(80) The study reported data from 218 individuals included in a test set (median age 51 years) enrolled from December 2019 to July 2021. Thirty-eight patients were female and 17% were Black or African American; other race or ethnicity data were not reported. The mean post-transplant time was 1,439 days (3.9 years). The study used a training set (n=149) to identify optimal % dd-cfDNA ($\geq 1\%$) and absolute values cut-offs ($\geq 78\text{cp/mL}$). Accuracy of dd-cfDNA testing was compared with the Molecular Microscope Diagnostic System (MMDx) and histological analysis using Banff criteria as reference standards. The use of two reference standards in this study is based on previous Trifecta analysis that suggested a strong correlation between dd-cfDNA fraction and molecular changes due to rejection assessed using MMDx.(81)

Table 7. Diagnostic Performance Characteristics of dd-cfDNA Tests for Detection of Active Kidney Allograft Rejection

Study; dd-cfDNA threshold	Biopsy-Matched Samples	Prevalence, n (%) ^a	Sensitivity, % (95% CI)	Specificity, % (95% CI)	AUC (95% CI)	PPV, % (95% CI)	NPV, % (95% CI)
Allosure							
Bloom et al (2017) ($\geq 1\%$)							
For-cause, dd-cfDNA	107	27 (25.2)	59 (44 to 74)	85 (79 to 81)	0.74 (0.61 to 0.86)	61 (NR)	84 (NR)
For-cause, SCr	204	58 (28.4)	NR	NR	0.54 (0.43 to 0.66)	NR	NR
Huang et al (2019) ($\geq 0.74\%$)							
For-cause, any rejection	63 (patients)	34 (54)	79.4 (NR)	72.4 (NR)	0.71 (0.58 to 0.85)	77.1 (NR)	75 (NR)
For-cause, CMR	63 (patients)	10 (16)	NR	NR	0.42 (0.17 to 0.66)	NR	NR
Prospera							
Sigdel et al (2019) ($\geq 1\%$)							
Overall, dd-cfDNA	217	33 (17.5)	88.7 (77.7 to 99.8)	72.6 (65.4 to 79.8)	0.87 (0.80 to 0.95)	52.0 (44.7 to 59.2) ^c	95.1 (90.5 to 99.7) ^c
Overall, eGFR	217	33 (17.5)	67.8 (51.3 to 84.2)	65.3 (57.6 to 73.0)	0.74 (0.66)	39.4 (31.6 to	85.9 (75.9 to

					to 0.83)	47.3) ^c	92.2) ^c
Surveillance, dd-cfDNA	114	12 (11.4)	92.3 (64.0 to 99.8)	75.2 (65.7 to 83.3)	0.89 (0.79 to 0.99)	55.4 (46.2 to 64.7) ^c 32.4 (24.8 to 41.1) ^d	96.7 (90.6 to 99.9) ^c 98.7 (92.0 to 99.8) ^d
For-cause, dd-cfDNA	103	25 (24.3)	84.0 (63.9 to 95.5) ^b	68.0 (56.4 to 78.1) ^b	NR	45.7 (36.8 to 54.8) ^d	93.0 (84.2 to 97.1) ^d
Halloran et al (2022) ≥1%; ≥78 cp/mL							
dd-cfDNA % + absolute quantity; MMDx criteria	218	71 (32.6)	83.1% (95% CI NR)	81.0% (95% CI NR)	0.88 (95% CI NR)	67.8% (95% CI NR)	90.8% (95% CI NR)
dd-cfDNA % + absolute quantity; Banff criteria	213	83 (39.0)	73.5% (95% CI NR)	80.8% (95% CI NR)	0.82 (95% CI NR)	70.9% (95% CI NR)	82.7% (95% CI NR)

AUC: area under the receiver-operating curve; CI: confidence interval; CMR: cell-mediated rejection; dd-cfDNA: donor-derived cell-free DNA; eGFR: estimated glomerular filtration rate; MMDx: molecular microscope diagnostic system; NPV: negative predictive value; NR: not reported; PPV: positive predictive value; SCr: serum creatinine.

^a Study disease prevalence.

^b Calculated based on reported case numbers.

^c Projected value as reported based on assumed disease prevalence of 25% in an at-risk population.

^d Calculated value based on study disease prevalence

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from randomized controlled trials.

No RCTs assessing the clinical utility of dd-cfDNA (i.e., AlloSure, Prospera) testing for renal allograft rejection were identified.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Because the clinical validity of dd-cfDNA (i.e., AlloSure, Prospera) testing to assess renal allograft rejection has not been established, a chain of evidence support clinical utility cannot be constructed.

Section Summary: Donor-Derived Cell-Free DNA Testing for Renal Transplant

Two meta-analyses were identified which pooled observational data from 9 studies. One reported an pooled sensitivity to detect graft rejection after kidney transplant of 70% with a specificity of 78%, and the other reported that antibody-mediated rejection was significantly associated with higher weighted median differences in dd-cfDNA fraction than in patients without rejection or in stable patients. Seven studies of the AlloSure test, using dd-cfDNA threshold values from $\geq 0.5\%$ to $\geq 1\%$, established a range of sensitivities from 59% to 86% with specificities of 72% to 100%. This corresponded to PPVs ranging from 61% to 77% and NPVs from 75% to 84%. Three studies provided information on the area under the curve of AlloSure to detect graft rejection which ranged from .8 to .996. A retrospective study of the Prospera test reported a PPV and NPV of 52% and 95%, respectively, using the $\geq 1\%$ dd-cfDNA threshold. A second, prospective Prospera study reported PPVs of 68% and 71% and NPVs 91% and 83% using combined dd-cfDNA fraction and absolute quantity compared with two different reference standards. Larger prospective studies validating dd-cfDNA thresholds for active rejection are needed to develop conclusions for each test. At present, no studies evaluating the clinical utility for AlloSure or Prospera dd-cfDNA testing were identified.

DONOR-DERIVED CELL-FREE DNA TESTING FOR LUNG TRANSPLANT

Clinical Context and Test Purpose

The purpose of dd-cfDNA testing in individuals with lung transplant who are undergoing surveillance is to detect allograft rejection.

The following PICO was used to select literature to inform this review.

Populations

The relevant population of interest are individuals with lung transplants who are undergoing surveillance for allograft rejection.

Interventions

The test being considered is dd-cfDNA testing to assess for lung allograft rejection.

A regular testing schedule is recommended for patients undergoing surveillance, with monthly testing in the first-year post-transplant and quarterly in the years 2-3. The proportion of dd-cfDNA relative to total cfDNA is reported. The report also notes that a threshold of $>0.85\%$ dd-cfDNA is associated with a higher probability of ACR, chronic lung allograft dysfunction (CLAD), and AMR and that the NPV is maximized at a $\%$ dd-cfDNA cutoff of 0.20%.

Comparators

The following test is currently being used to confirm a clinical suspicion of allograft rejection: bronchoscopy with transbronchial biopsy.

Outcomes

The general outcomes of interest are OS, test validity, morbid events, and hospitalizations. Follow-up over months to years is needed to monitor for signs of allograft rejection.

Beneficial outcomes resulting from a true-negative test result are avoiding unnecessary subsequent biopsy. Harmful outcomes resulting from a false-positive result may include an unnecessary biopsy or unnecessary treatment. Harmful outcomes from a false-negative result are increased risk of adverse transplant outcomes.

In a triage scenario, the test would need to identify precisely a group of patients that could safely forgo biopsy; therefore, the sensitivity, NPV, and negative likelihood ratio are key test performance characteristics.

Clinically Valid

A test must detect the presence or absence of a condition, the risk of developing a condition in the future, or treatment response (beneficial or adverse).

Study Selection Criteria

For the evaluation of clinical validity of dd-cfDNA testing, studies that met the criteria described in the first indication above.

Observational Studies

AlloSure

Sayah et al (2020) conducted a pilot study investigating the ability of AlloSure dd-cfDNA testing to detect ACR.(89) Biopsy-matched biorepository samples from 69 lung transplant recipients who had previously enrolled in the multicenter Lung Allograft Gene Expression Observational (LARGO) Study were evaluated. Diagnostic cohorts included patients with respiratory allograft infection (n=26), normal histopathology without infection or rejection (n=30), and ACR without concurrent infection (n=13). Samples were obtained between > 14 days and < 1-year post-transplant, and samples associated with potential concurrent infection with rejection were excluded. Median dd-cfDNA levels were 0.485% (IQR, 0.220 to 0.790) in the normal cohort, 1.52% (IQR, 0.520 to 2.550) in the ACR cohort, and 0.595% (IQR, 0.270 to 1.170) in the infection cohort. While dd-cfDNA levels were significantly higher in the ACR cohort compared to the normal cohort (p =.026), samples associated with infection were not significantly different from the normal (p =.282) or ACR (p=.1) cohorts. The AUC for detection of ACR was 0.717 (95% CI, 0.547 to 0.887; p=.025). At a threshold of 0.87% dd-cfDNA and an estimated prevalence rate of 25%, sensitivity for ACR was 73.1% (95% CI, 52.2% to 88.4%), specificity was 52.9% (95% CI, 27.8% to 77.0%), positive likelihood ratio was 1.55, negative likelihood ratio was 0.51, PPV was 34.1%, and NPV was 85.5%. The study is limited by the small sample size and use of archived samples, and raises concerns regarding the ability of AlloSure dd-cfDNA testing to detect AMR and to discriminate between infection and rejection.

Khush et al (2021) evaluated 107 biorepository plasma samples from 38 lung transplant recipients enrolled in the Genome Transplant Dynamics Study via AlloSure dd-cfDNA testing.(90) The study cohort included 14 patients (22 samples) with ACR confirmed by histopathology, 6 patients (7 samples) treated for ACR without a confirmed histopathological diagnosis, 6 patients (8 samples) with obstructive CLAD, 7 patients (9 samples) with AMR , 22 patients (33 samples) with infection without rejection, and 18 patients (28 samples) with stable allografts. The median dd-cfDNA levels in the ACR (0.91%; IQR, 0.39% to 2.07%) and CLAD (2.06%; IQR, 0.97% to 3.34%) cohorts were significantly higher compared to the stable cohort (p =.02, respectively). However, the AMR cohort was not statistically different when compared with the stable cohort (p =.07). The median dd-cfDNA level in an aggregated rejection cohort,

composed of ACR, AMR, and CLAD samples, was approximately 3-fold higher when compared to the stable cohort (1.06% versus 0.38%). At a dd-cfDNA threshold of 0.85%, the sensitivity for this spectrum of rejection was 55.6%, specificity was 75.8%, PPV was 43.3%, and NPV was 83.6%. The study is limited by the small sample size and use of archived samples. The authors suggest that AlloSure dd-cfDNA testing may have clinical utility as a plasma marker of "tissue injury" and that the 0.85% dd-cfDNA threshold requires further prospective clinical validation.

A retrospective study conducted by Keller et al (2022) included 157 patients enrolled in a post-transplant home surveillance program that included the AlloSure test for detection of acute allograft rejection.(91) The study analyzed data from patients at 4 U.S. centers. Data were collected from March to September 2020, during the COVID-19 pandemic at a time when in-office visits were limited and routine, surveillance bronchoscopy was deferred. Home monitoring was intended to identify those patients most at risk for acute rejection for triage to bronchoscopy. Study inclusion was limited to adults >18 years between 30 days and 3 years post-transplant. Of the total cohort, the mean age was 59 years and the majority were male (54%) and White race (64%). Eighteen percent were Black, 3% Asian, and 15% other race/ethnicity. The mean time since transplantation was 13 months, and 82% underwent bilateral transplantation. Diagnosis of ACR, AMR, infection, or a composite of these outcomes (Acute Lung Allograft Dysfunction [ALAD]), was made based on biopsy and/or clinical diagnosis. Mean dd-cfDNA % was 1.6% for acute rejection (ACR+AMR) and 1.7% for ALAD. In comparison, the mean dd-cfDNA in stable patients was 0.37%. Using a dd-cfDNA cut-off of 1.0% for detection of ALAD, the sensitivity was 73.9%, specificity 87.7%, PPV 43.4% and NPV 96.5%. Of the 157 patients with dd-cfDNA measurement for surveillance, 52 also had a contemporaneous reference standard surveillance bronchoscopy independent of dd-cfDNA level (i.e. patients who were not triaged to bronchoscopy). When analysis was limited to this subgroup, diagnostic performance declined slightly: 76.2% sensitivity, 70.0% specificity, 66.7% PPV and 79.2% NPV. The study was limited by the small sample size, particularly the limited number of unselected patients who underwent both dd-cfDNA testing and bronchoscopy.

Prospera Lung

Rosenheck et al (2022) assessed the predictive ability of dd-cfDNA testing using the Prospera test for lung transplant rejection.(92) The study included 195 samples from 103 patients, who were predominantly White (93%) and male (60%); mean age was 62 years. Black and Hispanic patients comprised 6% and 1% of the study population, respectively. The median time since lung transplant was 198 days, and most patients (85%) underwent lung biopsy for routine transplant surveillance. Consistent with other dd-cfDNA studies, median dd-cfDNA % was higher in patients with acute rejection (AR), which included ACR (1.43%) or AMR (2.50%), than those who were stable (0.46%). Prevalence of acute rejection was 28% (29/103), and prevalence of CLAD or neutrophilic-responsive allograft dysfunction (NRAD) was 21% (22/103); patients could be included in both diagnostic groups. Using a dd-cfDNA threshold of $\geq 1\%$ for prediction of acute rejection, sensitivity was 89.1% and specificity was 82.9%, resulting in an AUC of 0.91 (95% CI 0.83 to 0.98). PPV was 51.9% and NPV was 97.3%. For a combined measure that included AR, CLAD/NRAD, and infection, sensitivity was 59.9%, specificity 83.9%, AUC 0.76, PPV 43.6%, and NPV 91.0%. As with other dd-cfDNA studies in lung transplantation, this study was limited by the small sample size though unlike other studies samples were collected prospectively.

Clinically Useful

A test is clinically useful if the use of the results informs management decisions that improve the net health outcome of care. The net health outcome can be improved if patients receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

Direct Evidence

Direct evidence of clinical utility is provided by studies that have compared health outcomes for patients managed with and without the test. Because these are intervention studies, the preferred evidence would be from RCTs.

No RCTs assessing the clinical utility of dd-cfDNA (i.e., AlloSure or Prospera) testing to diagnose lung allograft rejection were identified.

Chain of Evidence

Indirect evidence on clinical utility rests on clinical validity. If the evidence is insufficient to demonstrate test performance, no inferences can be made about clinical utility.

Because the clinical validity of dd-cfDNA testing to assess for lung allograft rejection has not been established, a chain of evidence to support clinical utility cannot be constructed.

Section Summary: Donor-Derived Cell-Free DNA Testing for Lung Transplant

Four small diagnostic accuracy studies of dd-cfDNA testing with AlloSure or Prospera utilizing biorepository (3 studies) or prospectively collected samples (1 study) were identified. At a threshold of 0.87% dd-cfDNA, the PPV and NPV for detecting ACR in the first study were 34.1% and 85.5%, respectively. A second study reported a PPV of 43.3% and NPV of 83.6% at a dd-cfDNA cutoff of 0.85% for an aggregate rejection cohort composed of patients with ACR, AMR, and CLAD. In the third study, using a dd-cfDNA cut-off of 1.0%, PPV was 51.9% and NPV was 97.3% for acute rejection, and 43.6%, and 91.0% for acute rejection, CLAD/NRAD or infection. One study that used dd-cfDNA testing as part of a home surveillance program found a PPV 43.4% and NPV 96.5% for detection of ACR, AMR or infection, but when limited to patients with a contemporaneous reference standard surveillance bronchoscopy independent of dd-cfDNA level PPV 66.7% and NPV was 79.2%. Larger and additional prospective studies validating the dd-cfDNA threshold for active rejection are needed to develop conclusions. At present, no studies evaluating the clinical utility for AlloSure or Prospera dd-cfDNA testing were identified.

SUMMARY OF EVIDENCE

For individuals who have chronic heart failure who receive the sST2 assay to determine prognosis and/or to guide management, the evidence includes correlational studies and 2 meta-analyses. Relevant outcomes are overall survival, quality of life, and hospitalization. Most of the evidence is from reanalysis of existing randomized controlled trials and not from studies specifically designed to evaluate the predictive accuracy of sST2, and prospective and retrospective cross-sectional studies made up a large part of 1 meta-analysis. Studies have mainly found that elevated sST2 levels are statistically associated with elevated risk of mortality. A pooled analysis of study results found that sST2 significantly predicted overall mortality and cardiovascular mortality. Several studies, however, found that sST2 test results did not provide additional prognostic information compared with N-terminal pro B-type natriuretic peptide levels. Moreover, no comparative studies were identified on the use of the

sST2 assay to guide management of patients diagnosed with chronic heart failure. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have heart transplantation who receive sST2 assay to determine prognosis and/or to predict acute cellular rejection, the evidence includes a small number of retrospective observational studies on the Presage ST2 Assay. Relevant outcomes are overall survival, morbid events, and hospitalization. No prospective studies were identified that provide high-quality evidence on the ability of sST2 to predict transplant outcomes. One retrospective study (n = 241) found that sST2 levels were associated with acute cellular rejection and mortality; another study (n = 26) found that sST2 levels were higher during an acute rejection episode than before rejection. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have heart transplantation who receive peripheral blood measurement of donor derived cell free DNA assays to determine acute cellular rejection, the evidence includes observational studies. A validation study using 158 matched endomyocardial biopsy-plasma pairs from 76 pediatric and adult heart transplant recipients (ages 2 months or older, and 8 days more post-transplant) found a donor-specific fraction cutoff (0.32%) that produced a 100% negative predictive value for Grade 2 or higher acute cellular rejection. A prospective observational blinded study (n=174; pediatric=101, adult=73) using biopsy-paired samples found that myTAI_{HEART} level was associated with acute cellular and antibody-mediated rejection in both adult and pediatric heart transplant populations, and that an optimal donor fraction threshold (0.3%) ruled out the presence of either acute cellular rejection or antibody-mediated rejection. Both studies received industry funding. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have a heart transplant who receive a measurement of volatile organic compounds to assess cardiac allograft rejection, the evidence includes a diagnostic accuracy study. The relevant outcomes are overall survival, test validity, morbid events, and hospitalizations. The published study found that, for identifying grade 3 (now grade 2R) rejection, the negative predictive value of the breath test the study evaluated (97.2%) was similar to endomyocardial biopsy (96.7%) and the sensitivity of the breath test 78.6% was better than that for biopsy (42.4%). However, the breath test had lower specificity (62.4%) and a lower positive predictive value (5.6%) in assessing grade 3 rejection than biopsy (specificity, 97%; positive predictive value, 45.2%). The breath test was also not evaluated for grade 4 rejection. This single study is not sufficient to determine the clinical validity of the test measuring volatile organic compounds and no studies on clinical utility were identified. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with a heart transplant who receive gene expression profiling (GEP) to assess cardiac allograft rejection, the evidence includes 2 diagnostic accuracy studies and several randomized controlled trials (RCTs) evaluating clinical utility. The relevant outcomes are overall survival, test validity, morbid events, and hospitalizations. The 2 studies (CARGO, CARGO II) examining the diagnostic performance of GEP for detecting moderate-to-severe rejection lack of a consistent threshold for defining a positive GEP test (i.e., 20, 30, or 34) for determining positivity and a small number of positive cases. In the available studies, although the negative predictive values were relatively high (i.e., at least 88%), the performance

characteristics were only calculated based on few cases of rejection; therefore, performance data may be imprecise. Moreover, the positive predictive value in CARGO II was only 4.0% for patients who were at least 2 to 6 months posttransplant and 4.3% for patients more than 6 months posttransplant. The threshold indicating a positive test that seems to be currently accepted (a score of 34) was not prespecified; rather it evolved partway through the data collection period in the IMAGE study. In addition, the IMAGE study had several methodologic limitations (e.g., lack of blinding); further, the IMAGE study failed to provide evidence that GEP offers incremental benefit over biopsy performed on the basis of clinical exam or echocardiography. Patients at the highest risk of transplant rejection are patients within one year of the transplant, and, for that subset, there remains insufficient data on which to evaluate the clinical utility of GEP. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with a heart transplant who receive the combination test of gene expression profiling (GEP) with testing of dd-cfDNA to assess cardiac allograft rejection the evidence includes multiple studies and articles which are affiliated with the manufacturer. The relevant outcomes are overall survival, test validity, morbid events, and hospitalizations. At this time there is a lack of objective literature comparing dd-cfDNA to the current standard of care. Clinical trials are underway. Published studies either report (1) conflicts of interest, (2) lack control groups and/or (3) have serious methodologic problems that prevent the drawing of treatment-guiding conclusions. Peer reviewed, well-designed, well conducted, objective studies which indicate the technology can directly improve net health outcomes are needed.

For individuals with a renal transplant who are undergoing surveillance or have clinical suspicion of allograft rejection who receive testing of dd-cfDNA to assess renal allograft rejection, the evidence includes small diagnostic accuracy studies. Relevant outcomes are OS, test validity, morbid events, and hospitalizations. Two studies examined the diagnostic performance of dd-cfDNA for detecting moderate-to-severe rejection; the NPV was moderately high (75% to 84%), and performance characteristics were calculated on 27 cases of active transplant rejection. The threshold indicating a positive test was not prespecified. A subsequent smaller single-center study that explored variation in clinical validity based on different rejection mechanisms found the strongest performance characteristics for AlloSure with antibody-mediated rejection. Using dd-cfDNA threshold values from $\geq 0.5\%$ to $\geq 1\%$, the Allosure test established a range of sensitivities from 59% to 86% and specificities of 72% to 100% for the detection of graft rejection. This corresponded to PPVs ranging from 61% to 77% and NPVs from 75% to 84%. A retrospective single-center study of the Prospera dd-cfDNA test reported a PPV and NPV of 52% and 95%, respectively, for detection of active rejection among a combined cohort of patients undergoing surveillance or for-cause biopsies, using the 1% dd-cfDNA threshold previously proposed for the AlloSure test. Larger prospective studies validating the dd-cfDNA thresholds for active rejection are needed to develop conclusions for each test. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with a lung transplant who receive testing of dd-cfDNA to assess lung allograft rejection, the evidence includes 4 small diagnostic accuracy studies. Relevant outcomes are OS, test validity, morbid events, and hospitalizations. One study examined the diagnostic performance of dd-cfDNA testing at a threshold of 0.87% for detecting acute cellular rejection, yielding a PPV of 34.1% and a NPV of 85.5%. A second study reported a PPV of 43.3% and NPV of 83.6% for an aggregate rejection cohort composed of patients with acute cellular

rejection, antibody-mediated rejection, and CLAD. In the third study, using a dd-cfDNA cut-off of 1.0%, PPV was 51.9% and NPV was 97.3% for acute rejection, and 43.6%, and 91.0% for acute rejection, CLAD/NRAD or infection. One study that used dd-cfDNA testing as part of a home surveillance program found a PPV 43.4% and NPV 96.5% for detection of ACR, AMR or infection, but when limited to patients with a contemporaneous reference standard surveillance bronchoscopy independent of dd-cfDNA level, PPV 66.7% and NPV was 79.2%. All 4 studies were limited by small sample sizes, and no clinical utility studies were identified. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Supplemental Information

CLINICAL INPUT RECEIVED THROUGH PHYSICIAN SPECIALTY SOCIETIES AND ACADEMIC MEDICAL CENTERS

While the various physician specialty societies and academic medical centers may collaborate with and make recommendations during this process, through the provision of appropriate reviewers, input received does not represent an endorsement or position statement by the physician specialty societies or academic medical centers, unless otherwise noted.

2012 Input

In response to requests, Blue Cross Blue Shield Association received input from 7 academic medical centers and 1 specialty society while this policy was under review in December 2011. The input was mixed on the question of whether Allomap should be investigational. Four reviewers agreed with the investigational status, 1 disagreed, and 3 indicated it was a split decision/other. The reviewers were generally in agreement that the sensitivity and specificity has not yet been adequately defined for Allomap and that the NPV was not sufficiently high to preclude the need for biopsy. There was mixed input about the need for surveillance cardiac biopsies to be performed in the absence of clinical signs and/or symptoms of rejection.

2008 Input

In response to requests, Blue Cross Blue Shield Association received input from 2 academic medical centers and 2 physician specialty societies while this policy was under review in 2008. Three reviewers agreed that these approaches for monitoring heart transplant rejection are considered investigational. The American College of Cardiology (ACC) disagreed with the policy, stating that the American College of Cardiology disagreed with the policy, stating that the College considers the available laboratory tests to have good potential to diagnose heart transplant rejection and reduce the frequency of invasive biopsies performed on heart transplant patients, although questions remained as to their role in clinical practice.

PRACTICE GUIDELINES AND POSITION STATEMENTS

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American College of Cardiology et al

In 2022, the American College of Cardiology, American Heart Association, and Heart Failure Society issued updated an guideline for the management of heart failure.(8) The 2022 guideline replaced a 2013 guideline (1) and a 2017 focused guideline update. (98) The guideline states measurement of natriuretic peptide levels may be useful for diagnosis, risk stratification, and prognosis of heart failure. The use of soluble suppression oftumorigenicity-2 is not discussed specifically, though the guideline notes that "a widening array of biomarkers including markers of myocardial injury, inflammation, oxidative stress, vascular dysfunction, and matrix remodeling have been shown to provide incremental prognostic information over natriuretic peptides but remain without evidence of an incremental management benefit."

American Society of Transplant Surgeons

In 2023, the American Society of Transplant Surgeons (ASTS) issued the following position statements applicable to adult populations.(99)

- We suggest that clinicians consider measuring serial dd-cfDNA levels in kidney transplant recipients with stable renal allograft function to exclude the presence of subclinical antibody-mediated rejection.
- We recommend that clinicians measure dd-cfDNA levels in kidney transplant recipients with acute allograft dysfunction to exclude the presence of rejection, particularly antibody-mediated rejection (ABMR).
- We do not recommend the use of blood gene expression profiling (GEP) in kidney transplant recipients for the purpose of diagnosing or excluding sub-clinical rejection, as adequate evidence supporting such use is still lacking.
- We do not recommend the use of blood GEP to diagnose or exclude the presence of acute graft rejection in kidney transplant recipients with acute allograft dysfunction given the paucity of data to support this practice.
- We recommend that dd-cfDNA may be utilized to rule out subclinical rejection in heart transplant recipients.
- We recommend that clinicians utilize peripheral blood GEP as a non-invasive diagnostic tool to rule out acute cellular rejection in stable, low-risk, adult heart transplant recipients who are over 55 days status post heart transplantation."
- We recommend that there is still insufficient evidence to recommend dd-cfDNA or GEP testing in liver transplant recipients.

"Caveats and recommendations for future studies:

- None of these recommendations should be construed as recommending one biomarker over another in the same diagnostic niche.
- We strongly recommend ongoing clinical studies to clarify the scenarios in which molecular diagnostic studies should be utilized.
- We specifically recommend that studies be carried out to evaluate the potential role of dd-cfDNA surveillance in kidney transplant recipients to improve long-term allograft survival."

International Society of Heart and Lung Transplantation

In 2022, the International Society of Heart and Lung Transplantation issued updated guidelines for the care of heart transplant recipients.(100) The guidelines included the following recommendations (see Table 8).

Table 8. Guidelines for Postoperative Care of Heart Transplant Recipients

Recommendation	COR	LOE
"It is reasonable to perform periodic EMB during the first 3 to 12 postoperative months for surveillance of HT rejection."	Ila	C
"After the first post-operative year, it is reasonable to continue EMB surveillance in patients who are at higher risk for late acute rejection..."	Ila	C
"Gene Expression Profiling (GEP) (i.e., AlloMap) of peripheral blood can be used in low-risk patients between 2 months and 5 years after HT to identify adult recipients who have low risk of current ACR to reduce the frequency of EMB. Data in children does not allow a general recommendation of GEP as a routine tool at present."	Ila	B

ACR: acute cellular rejection; COR: class of recommendation; EMB: endomyocardial biopsy; HT: heart transplant; LOE: level of evidence.

American Society of Transplantation (AST)

The AST's Thoracic and Critical Care Community convened a virtual expert panel in heart and kidney transplantation to evaluate emerging diagnostics and how they may be best utilized to monitor and manage transplant patients on January 27-28, 2022. In regard to the evolving use of biomarkers in heart transplantation, there was broad agreement that larger and more sophisticated studies are needed to investigate, compare, and contrast the growing number of potential biomarkers in heart transplant medicine. In addition, it is clear that the endomyocardial biopsy is not the gold standard to detect rejection and that we need a better definition of rejection to validate all current and future biomarkers. There was also recognition that there is unlikely to exist a single, preeminent assay or biomarker which serves all purposes, and that combinations of biomarkers, employed either together or at different times after transplant, is the most probable future outcome. 71% of the members of the expert panel agreed that multicenter clinical trials are necessary to support regulatory endorsement and the routine clinical use of a biomarker/diagnostic test. (Kobashigawa et al., 2023). (103)

Kidney Disease Improving Global Outcomes

The Kidney Disease Improving Global Outcomes (2009) issued guidelines for the care of kidney transplant recipients.(104) The guidelines included the following recommendations (see Table 9).

Table 9. Guidelines for Biopsy in Renal Transplant Recipients

Recommendation	SOR	LOE
"We recommend kidney allograft biopsy when there is a persistent, unexplained increase in serum creatinine."	Level 1	C
"We suggest kidney allograft biopsy when serum creatinine has not returned to baseline after treatment of acute rejection."	Level 2	D
"We suggest kidney allograft biopsy every 7-10 days during delayed function."	Level 2	C
"We suggest kidney allograft biopsy if expected kidney function is not achieved within the first 1-2 months after transplantation."	Level 2	D
"We suggest kidney allograft biopsy when there is new onset of proteinuria."	Level 2	C
"We suggest kidney allograft biopsy when there is unexplained proteinuria ≥ 3.0 g/g creatinine or ≥ 3.0 g per 24 hours."	Level 2	C

LOE: level of evidence; SOR: strength of recommendation.

KDGO does not list any gene expression or cell-free DNA techniques in their guidelines for managing transplant recipients.

European Association of Urology

The European Association of Urology (2018) posted guidelines which indicate that the ultimate standard for the diagnosis of rejection is transplant biopsy.(105) It is impossible to differentiate acute rejection solely on clinical indicators from other causes of renal dysfunction (e.g., acute tubular necrosis, infection, disease recurrence or CNI nephrotoxicity). Therefore, all rejections should be verified by renal biopsy.

EAU does not list any gene expression or cell-free DNA techniques in their guidelines for managing transplant recipients.

U.S. PREVENTIVE SERVICES TASK FORCE RECOMMENDATIONS

Not applicable

ONGOING AND UNPUBLISHED CLINICAL TRIALS

Some currently ongoing and unpublished trials that might influence this review are listed in Table 10.

Table 10. Summary of Key Active Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
AlloMap			
Ongoing			
NCT01833195 ^a	Outcomes AlloMap Registry: the Long-term Management and Outcomes of Heart Transplant Recipients With AlloMap Testing (OAR)	2444	Feb 2020 (active, not recruiting)
NCT02178943 ^a	Utility of Donor-Derived Cell-free DNA in Association With Gene-Expression Profiling (AlloMap®) in Heart Transplant Recipients (D-OAR)	100	Feb 2020 (unknown)
HeartCare			
Ongoing			
NCT03695601 ^a	Surveillance HeartCare Outcomes Registry (SHORE)	3450	Jun 2027 (recruiting)
AlloSure (Kidney)			
Ongoing			
NCT04566055 ^a	Assessing AlloSure dd-cfDNA Monitoring Insights of Renal Allografts With Longitudinal Surveillance (ADMIRAL)	1000	Oct 2020 (active, not recruiting)
NCT04057742 ^a	AlloSure for the Monitoring of Antibody Mediated Processes After Kidney Transplantation (All-MAP)	69	Dec 2024 (recruiting)
NCT03326076 ^a	Evaluation of Patient Outcomes From the Kidney Allograft Outcomes AlloSure Registry (KOAR)	4000	Dec 2025 (recruiting)
NCT04601155 ^a	Transition of Renal Patients Using AlloSure Into Community Kidney Care (TRACK)	3500	Sep 2026 (recruiting)
AlloSure (Lung)			

Ongoing			
NCT04318587 ^a	Assessment of Donor Derived Cell Free DNA and Utility in Lung Transplantation	50	Sep 2023 (recruiting)
NCT05050955 ^a	Allosure Lung Assessment and Metagenomics Outcomes Study (ALAMO)	1500	Dec 2026 (not yet recruiting)
Prospera (Kidney) Ongoing			
NCT04239703 ^a	INTERCOMEX DD-cfDNA-HLA-MMDx Study: Comparing the DD-cfDNA Test to MMDx Microarray Test, Central HLA Antibody Testx, and Histology	300	Dec 2022 (recruiting)
NCT04091984 ^a	The PROspera Kidney Transplant ACTIVE Rejection Assessment Registry (ProActive)	3000	Oct 2027 (recruiting)
NCT03984747 ^a	Study for the Prediction of Active Rejection in Organs Using Donor-derived Cell-free DNA Detection (SPARO)	500	Oct 2028 (recruiting)
Prospera (Heart) Ongoing			
NCT04707872 ^a	Trifecta-Heart cfDNA-MMDx Study: Comparing the DD-cfDNA test to MMDx Microarray Test and Central HLA Antibody Test	300	Dec 2022 (recruiting)
NCT05081739 ^a	Donor-Derived Cell-free DNA to Detect Rejection in Cardiac Transplantation (DETECT)	600	Jan 2025 (not yet recruiting)
NCT05205551	Prospera Test Evaluation in Cardiac Transplant (ProTECT)	1000	Dec 2027 (recruiting)
Prospera (Lung) Ongoing			
NCT05837663 ^a	Trifecta-Lung cfDNA-MMDx Study: Comparing the Dd-cfDNA Test to MMDx Microarray Test and Central HLA Antibody Test	600	Dec 2025 (recruiting)
NCT05170425 ^a	Observational Registry Study With Sub-analysis (Patients Previously Randomized to LAMBDA 001) to Assess Prospera TM Performance for Detection of CLAD After Lung Transplant (LAMBDA 002)	1000	Dec 2029 (not yet recruiting)

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

Government Regulations

National:

Heartsbreath Test for Heart Transplant Rejection; Pub number: 100.3; Version 1; Section Number 260.10; *Effective date:12/8/08*

The Centers for Medicare & Medicaid Services has determined that the evidence does not adequately define the technical characteristics of the test nor demonstrate that Heartsbreath testing to predict heart transplant rejection improves health outcomes in Medicare beneficiaries. Thus, we conclude that the Heartsbreath test is not reasonable and necessary under section 1862(a)(1)(A) of the Social Security Act and is non-covered.(106)

There is no national coverage determination on use of the **AlloMap** test after heart transplant.

There is no national coverage determination addressing **AlloSure**.

There is no **national or local coverage** determination for **ST2 assays**.

- The Medicare Clinical Diagnostic Laboratory Fee Schedule lists a fee for 83006.

Local:

WPS - MoIDX: Molecular Testing for Solid Organ Allograft Rejection (L38680)

For services performed on or after 06/06/2021; For services performed on or after 07/27/2023 (107)

Coverage Guidance

Molecular diagnostic tests that assess a transplanted allograft for rejection status are covered when ALL of the following criteria are met:

- The test must provide information about at least one of the two following clinical status determinations:
 - AR status
 - Cellular or Antibody-mediated rejection (ACR or AMR) status
- The intended use of the test must be:
 - To assist in the evaluation of adequacy of immunosuppression, wherein a non-invasive or minimally invasive test can be used in lieu of a tissue biopsy in a patient for whom information from a tissue biopsy would be used to make a management decision regarding immunosuppression, OR
 - As a rule-out test for AR in validated populations of patients with clinical suspicion of rejection with a non-invasive or minimally invasive test to make a clinical decision regarding obtaining a biopsy, OR
 - For further evaluation of allograft status for the probability of allograft rejection after a physician-assessed pretest, OR
 - To assess rejection status in patients that have received a biopsy, but the biopsy results are inconclusive or limited by insufficient material.
- The test demonstrates analytical validity (AV), including an analytical and clinical validation for any given measured analytes, and has demonstrated equivalence or superiority for sensitivity or specificity (depending on intended use) of detecting allograft rejection to other already-accepted tests for the same intended use measuring the same or directly comparable analytes.
- Clinical validity (CV) of any analytes (or expression profiles) measured must be established through a study published in the peer-reviewed literature for the intended use of the test in the intended population. The degree of validity must be similar or superior to established and covered tests (see associated coverage Articles). If conducted with concordance to tissue histologic evaluation the Banff Classification for renal allografts or other accepted criteria (if existing) for other organs must be used.
- The test is being used in a patient who is part of the population in which the test was analytically validated and has demonstrated CV.
- For a given patient encounter, only one molecular test for assessing allograft status may be performed UNLESS a second test, meeting all the criteria established herein, is reasonable and necessary as an adjunct to the first test.
- For minimally or non-invasive tests, the benefit to risk profile of the molecular test is considered by the ordering clinician to be more favorable than the benefit to risk profile

of a tissue biopsy, or a tissue biopsy cannot be obtained. For example, this may be the case if a biopsy is considered medically contraindicated in a patient.

- The test successfully completes a Technical Assessment that will ensure that AV, CV, and clinical utility criteria set in this policy are met to establish the test as Reasonable and Necessary.
- Covered tests with AV that is significantly below similar services may have coverage rescinded.

WPS – Billing and Coding: MolDX: Molecular Testing for Solid Organ Allograft Rejection (A58207)

Original Effective Date: 6/6/21; Revision Effective Date: 2/29/24 (108)

The information in this article contains billing, coding or other guidelines that complement the Local Coverage Determination (LCD) for MolDX: Molecular Testing for Solid Organ Allograft Rejection L38680.

The tests below have completed a MolDx technical assessment and are deemed compliant with policy. However, the tests must be billed in accordance with the specific required indications/coverage criteria outlined in the policy.

Test (Manufacturer)	Methodology	Specimen Source	Analyte(s) and Principle of Test	Transplant Type(s)
AlloMap® (CareDx)	GEP (qRT- PCR)	Blood	GEP is used to calculate a score associated with the level of immune quiescence and probability of AR	Heart
AlloSure® (CareDx)	NGS (targeting 266 SNPs)	Blood	% dd-cfDNA is measured to identify graft injury and AR	Kidney Heart Lung
HeartCare (CareDx)	AlloSure Heart + AlloMap Heart performed on the same patient sample	Blood	See AlloSure Heart and AlloMap Heart	Heart
Prospera™ (Natera™)	mmPCR-NGS (targeting 13,926 SNP's)	Blood	% dd-cfDNA is measured to identify graft injury and AR	Kidney Heart Lung
QSant™ (NephroSant)	Multianalyte assay with algorithmic analysis	Urine	The concentrations of 6 urinary biomarkers (cell-free DNA (cfDNA), methylated cfDNA (m-cfDNA), CXCL10, clusterin, total protein and creatinine) are incorporated into an algorithm to calculate a composite score	Kidney

			(Q-score), to demonstrate the probability of rejection risk.	
TruGraf® (Transplant Genomics)	GEP (RT-PCR)	Blood	GEP signatures identify patients that are adequately immunosuppressed	Kidney
Viracor TRAC™ (Transplant Rejection Allograft Check) (Eurofins)	Low-coverage WGS querying >100,000 SNPs	Blood	% dd-cfDNA is measured to identify AR	Kidney
VitaGraft/ Allaro Kidney (Oncocyte)	droplet digital PCR (ddPCR)	Blood	% dd-cfDNA is measured to identify graft injury and AR	Kidney

AR: acute rejection; dd-cfDNA: donor-derived cell-free DNA; GEP: gene expression profile; NGS: next generation sequencing; PCR: polymerase chain reaction; WGS: whole genome sequencing

(The above Medicare information is current as of the review date for this policy. However, the coverage issues and policies maintained by the Centers for Medicare & Medicare Services [CMS, formerly HCFA] are updated and/or revised periodically. Therefore, the most current CMS information may not be contained in this document. For the most current information, the reader should contact an official Medicare source.)

Related Policies

- Galectin-3 Testing in the Assessment and Management of Chronic Heart Failure
- Heart Transplant
- Heart/Kidney Transplant (Combined)
- Heart/Lung Transplant (Combined)
- Measurement of Lipoprotein-associated Phospholipase A2 (Lp-PLA2) and Secretory Type II Phospholipase A2 (sPLA2-IIA) and Secretory Type II Phospholipase A2 (sPLA2-IIA) in the Assessment of Cardiovascular Risk
- Novel Biomarkers in Risk Assessment and Management of Cardiovascular Disease
- Skin Advanced Glycation End Products (AGE) Measurement By Multi-Wavelength Fluorescent Spectroscopy

References

1. Yancy CW, Jessup M, Bozkurt B, et al. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. J Am Coll Cardiol. Oct 15 2013; 62(16): e147-239. PMID 23747642
2. Roger VL, Go AS, Lloyd-Jones DM, et al. Heart disease and stroke statistics--2011 update: a report from the American Heart Association. Circulation. Feb 01 2011; 123(4): e18-e209. PMID 21160056
3. Virani SS, Alonso A, Benjamin EJ, et al. Heart Disease and Stroke Statistics-2020 Update: A Report From the American Heart Association. Circulation. Mar 03 2020; 141(9): e139-e596. PMID 31992061

4. Rohde LE, Beck-da-Silva L, Goldraich L, et al. Reliability and prognostic value of traditional signs and symptoms in outpatients with congestive heart failure. *Can J Cardiol.* May 15 2004; 20(7): 697-702. PMID 15197422
5. Marcus GM, Gerber IL, McKeown BH, et al. Association between phonocardiographic third and fourth heart sounds and objective measures of left ventricular function. *JAMA.* May 11 2005; 293(18): 2238-44. PMID 15886379
6. Stevenson LW, Perloff JK. The limited reliability of physical signs for estimating hemodynamics in chronic heart failure. *JAMA.* Feb 10 1989; 261(6): 884-8. PMID 2913385
7. Gaggin HK, Januzzi JL. Biomarkers and diagnostics in heart failure. *Biochim Biophys Acta.* Dec 2013; 1832(12): 2442-50. PMID 23313577
8. Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation.* May 03 2022; 145(18): e895-e1032. PMID35363499
9. McMurray JJ, Adamopoulos S, Anker SD, et al. ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2012: The Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur J Heart Fail.* Aug 2012; 14(8): 803-69. PMID 22828712
10. Lindenfeld J, Albert NM, Boehmer JP, et al. HFSA 2010 Comprehensive Heart Failure Practice Guideline. *J Card Fail.* Jun 2010; 16(6): e1-194. PMID 20610207
11. Savarese G, Trimarco B, Dellegrottaglie S, et al. Natriuretic peptide-guided therapy in chronic heart failure: a meta-analysis of 2,686 patients in 12 randomized trials. *PLoS One.* 2013; 8(3): e58287. PMID 23472172
12. Troughton RW, Frampton CM, Brunner-La Rocca HP, et al. Effect of B-type natriuretic peptide-guided treatment of chronic heart failure on total mortality and hospitalization: an individual patient meta-analysis. *Eur Heart J.* Jun 14 2014; 35(23): 1559-67. PMID 24603309
13. Bhardwaj A, Januzzi JL. ST2: a novel biomarker for heart failure. *Expert Rev Mol Diagn.* May 2010; 10(4): 459-64. PMID 20465500
14. Chowdhury P, Kehl D, Choudhary R, et al. The use of biomarkers in the patient with heart failure. *Curr Cardiol Rep.* Jun 2013; 15(6): 372. PMID 23644993
15. Ciccone MM, Cortese F, Gesualdo M, et al. A novel cardiac bio-marker: ST2: a review. *Molecules.* Dec 11 2013; 18(12): 15314-28. PMID 24335613
16. Daniels LB, Bayes-Genis A. Using ST2 in cardiovascular patients: a review. *Future Cardiol.* Jul 2014; 10(4): 525-39. PMID 25301315
17. Dieplinger B, Mueller T. Soluble ST2 in heart failure. *Clin Chim Acta.* Mar 30 2015; 443: 57-70. PMID 25269091
18. Mueller T, Dieplinger B. The Presage((R)) ST2 Assay: analytical considerations and clinical applications for a high-sensitivity assay for measurement of soluble ST2. *Expert Rev Mol Diagn.* Jan 2013; 13(1): 13-30. PMID 23256700
19. Shah RV, Januzzi JL. ST2: a novel remodeling biomarker in acute and chronic heart failure. *Curr Heart Fail Rep.* Mar 2010; 7(1): 9-14. PMID 20425491
20. Xu D, Chan WL, Leung BP, et al. Selective expression of a stable cell surface molecule on type 2 but not type 1 helper T cells. *J Exp Med.* Mar 02 1998; 187(5): 787-94. PMID 9480988
21. Weinberg EO, Shimpo M, De Keulenaer GW, et al. Expression and regulation of ST2, an interleukin-1 receptor family member, in cardiomyocytes and myocardial infarction. *Circulation.* Dec 03 2002; 106(23): 2961-6. PMID 12460879

22. Stewart S, Winters GL, Fishbein MC, et al. Revision of the 1990 working formulation for the standardization of nomenclature in the diagnosis of heart rejection. *J Heart Lung Transplant*. Nov 2005; 24(11): 1710-20. PMID 16297770
23. Celec P, Vlkova B, Laukova L, et al. Cell-free DNA: the role in pathophysiology and as a biomarker in kidney diseases. *Expert Rev Mol Med*. Jan 18 2018; 20: e1. PMID 29343314
24. Natera. Prospera Heart Overview. 2022.
25. North PE, Ziegler E, Mahnke DK, et al. Cell-free DNA donor fraction analysis in pediatric and adult heart transplant patients by multiplexed allele-specific quantitative PCR: Validation of a rapid and highly sensitive clinical test for stratification of rejection probability. *PLoS One*. 2020; 15(1): e0227385. PMID 31929557
26. CareDx. Overview: AlloMap Testing: Answering Unmet Needs in Heart Transplant Surveillance. n.d.; <http://www.allomap.com/providers/overview/>.
27. Duong Van Huyen JP, Tible M, Gay A, et al. MicroRNAs as non-invasive biomarkers of heart transplant rejection. *Eur Heart J*. Dec 01 2014; 35(45): 3194-202. PMID 25176944
28. Patel PC, Hill DA, Ayers CR, et al. High-sensitivity cardiac troponin I assay to screen for acute rejection in patients with heart transplant. *Circ Heart Fail*. May 2014; 7(3): 463-9. PMID 24733367
29. Organ Procurement and Transplantation Network. National Data. 2018; <https://optn.transplant.hrsa.gov/data/view-data-reports/national-data/#>. Accessed August 31, 2020.
30. Hariharan S, Israni AK, Danovitch G. Long-Term Survival after Kidney Transplantation. *N Engl J Med*. Aug 19 2021; 385(8): 729-743. PMID 34407344
31. Goldberg RJ, Weng FL, Kandula P. Acute and Chronic Allograft Dysfunction in Kidney Transplant Recipients. *Med Clin North Am*. May 2016; 100(3): 487-503. PMID 27095641
32. Ahmad I. Biopsy of the transplanted kidney. *Semin Intervent Radiol*. Dec 2004; 21(4): 275-81. PMID 21331139
33. Solez K, Colvin RB, Racusen LC, et al. Banff 07 classification of renal allograft pathology: updates and future directions. *Am J Transplant*. Apr 2008; 8(4): 753-60. PMID 18294345
34. Haas M. The Revised (2013) Banff Classification for Antibody-Mediated Rejection of Renal Allografts: Update, Difficulties, and Future Considerations. *Am J Transplant*. May 2016; 16(5): 1352-7. PMID 26696524
35. Natera. Prospera Transplant Assessment: Indications for Use. 2020; <https://www.natera.com/wp-content/uploads/2020/11/Organ-Health-Clinician-Indications-for-use.pdf>.
36. Natera. Prospera Transplant Assessment: Clinician's Guide to Results. 2019; <https://www.natera.com/wp-content/uploads/2020/11/Organ-Health-Clinician-Clinicians-guide-to-results.pdf>.
37. Cohn JN, Tognoni G. A randomized trial of the angiotensin-receptor blocker valsartan in chronic heart failure. *N Engl J Med*. Dec 06 2001; 345(23): 1667-75. PMID 11759645
38. O'Connor CM, Whellan DJ, Lee KL, et al. Efficacy and safety of exercise training in patients with chronic heart failure: HF-ACTION randomized controlled trial. *JAMA*. Apr 08 2009; 301(14): 1439-50. PMID 19351941
39. Kjekshus J, Apetrei E, Barrios V, et al. Rosuvastatin in older patients with systolic heart failure. *N Engl J Med*. Nov 29 2007; 357(22): 2248-61. PMID 17984166
40. Januzzi JL, Rehman SU, Mohammed AA, et al. Use of amino-terminal pro-B-type natriuretic peptide to guide outpatient therapy of patients with chronic left ventricular systolic dysfunction. *J Am Coll Cardiol*. Oct 25 2011; 58(18): 1881-9. PMID 22018299
41. Ky B, French B, McCloskey K, et al. High-sensitivity ST2 for prediction of adverse outcomes in chronic heart failure. *Circ Heart Fail*. Mar 2011; 4(2): 180-7. PMID 21178018

42. Bayes-Genis A, de Antonio M, Galan A, et al. Combined use of high-sensitivity ST2 and NTproBNP to improve the prediction of death in heart failure. *Eur J Heart Fail.* Jan 2012; 14(1): 32-8. PMID 22179033
43. Broch K, Ueland T, Nymo SH, et al. Soluble ST2 is associated with adverse outcome in patients with heart failure of ischaemic aetiology. *Eur J Heart Fail.* Mar 2012; 14(3): 268-77. PMID 22302661
44. Felker GM, Fiuzat M, Thompson V, et al. Soluble ST2 in ambulatory patients with heart failure: Association with functional capacity and long-term outcomes. *Circ Heart Fail.* Nov 2013; 6(6): 1172-9. PMID 24103327
45. Gaggin HK, Motiwala S, Bhardwaj A, et al. Soluble concentrations of the interleukin receptor family member ST2 and -blocker therapy in chronic heart failure. *Circ Heart Fail.* Nov 2013; 6(6): 1206-13. PMID 24114865
46. Anand IS, Rector TS, Kuskowski M, et al. Prognostic value of soluble ST2 in the Valsartan Heart Failure Trial. *Circ Heart Fail.* May 2014; 7(3): 418-26. PMID 24622243
47. Zhang R, Zhang Y, An T, et al. Prognostic value of sST2 and galectin-3 for death relative to renal function in patients hospitalized for heart failure. *Biomark Med.* 2015; 9(5): 433-41. PMID 25985174
48. Dupuy AM, Curinier C, Kuster N, et al. Multi-Marker Strategy in Heart Failure: Combination of ST2 and CRP Predicts Poor Outcome. *PLoS One.* 2016; 11(6): e0157159. PMID 27311068
49. Aimo A, Vergaro G, Passino C, et al. Prognostic Value of Soluble Suppression of Tumorigenicity-2 in Chronic Heart Failure: A Meta-Analysis. *JACC Heart Fail.* Apr 2017; 5(4): 280-286. PMID 27816512
50. Januzzi JL, Horne BD, Moore SA, et al. Interleukin receptor family member ST2 concentrations in patients following heart transplantation. *Biomarkers.* May 2013; 18(3): 250-6. PMID 23557127
51. Pascual-Figal DA, Garrido IP, Blanco R, et al. Soluble ST2 is a marker for acute cardiac allograft rejection. *Ann Thorac Surg.* Dec 2011; 92(6): 2118-24. PMID 22035779
52. Phillips M, Boehmer JP, Cataneo RN, et al. Heart allograft rejection: detection with breath alkanes in low levels (the HARDBALL study). *J Heart Lung Transplant.* Jun 2004; 23(6): 701-8. PMID 15366430
53. Khush KK, Patel J, Pinney S, et al. Noninvasive detection of graft injury after heart transplant using donor-derived cell-free DNA: A prospective multicenter study. *Am J Transplant.* Oct 2019; 19(10): 2889-2899. PMID 30835940
54. Kim PJ, Olymbios M, Siu A, et al. A novel donor-derived cell-free DNA assay for the detection of acute rejection in heart transplantation. *J Heart Lung Transplant.* Jul 2022; 41(7): 919-927. PMID 35577713
55. Richmond ME, Zangwill SD, Kindel SJ, et al. Donor fraction cell-free DNA and rejection in adult and pediatric heart transplantation. *J Heart Lung Transplant.* May 2020; 39(5): 454-463. PMID 31983667
56. CareDx. HeartCare Interpretive Guide. April 2021; <https://caredx.com/wp-content/uploads/2021/09/LK-10482-Heartcare-Interpretive-Guide.pdf>.
57. Blue Cross Blue Shield Technology Evaluation Center (TEC). Gene expression profiling as a noninvasive method to monitor for cardiac allograft rejection. *TEC Assessment Program.* 2011;26(8).
58. Deng MC, Eisen HJ, Mehra MR, et al. Noninvasive discrimination of rejection in cardiac allograft recipients using gene expression profiling. *Am J Transplant.* Jan 2006; 6(1): 150-60. PMID 16433769

59. Crespo-Leiro MG, Stypmann J, Schulz U, et al. Clinical usefulness of gene-expression profile to rule out acute rejection after heart transplantation: CARGO II. *Eur Heart J*. Sep 01 2016; 37(33): 2591-601. PMID 26746629
60. Kanwar MK, Khush KK, Pinney S, et al. Impact of cytomegalovirus infection on gene expression profile in heart transplant recipients. *J Heart Lung Transplant*. Feb 2021; 40(2): 101-107. PMID 33341360
61. Gondi KT, Kao A, Linard J, et al. Single-center utilization of donor-derived cell-free DNA testing in the management of heart transplant patients. *Clin Transplant*. May 2021; 35(5): e14258. PMID 33606316
62. Kobashigawa J, Patel J, Azarbal B, et al. Randomized pilot trial of gene expression profiling versus heart biopsy in the first year after heart transplant: early invasive monitoring attenuation through gene expression trial. *Circ Heart Fail*. May 2015; 8(3): 557-64. PMID 25697852
63. Pham MX, Deng MC, Kfoury AG, et al. Molecular testing for long-term rejection surveillance in heart transplant recipients: design of the Invasive Monitoring Attenuation Through Gene Expression (IMAGE) trial. *J Heart Lung Transplant*. Aug 2007; 26(8): 808-14. PMID 17692784
64. Pham MX, Teuteberg JJ, Kfoury AG, et al. Gene-expression profiling for rejection surveillance after cardiac transplantation. *N Engl J Med*. May 20 2010; 362(20): 1890-900. PMID 20413602
65. CareDx Inc. HeartCare. n.d. <https://caredx.com/products-and-services/transplant-services/heart/heartcare/>. Accessed May 11, 2023
66. Kanwar M, Pinney S, Shah P, Kobashigawa J. Correlation of longitudinal gene-expression profiling (GEP) score to cytomegalovirus (CMV) infection: results from the outcomes Allomap® registry. *J Heart Lung Transpl*. 2020;39(4):S83-S84. doi:10.1016/j.healun.2020.01.1311
67. Purewal S, Moayedi Y, Runeckles K, et al. Are two tests better than one? Combining donor derived cell-free DNA and gene expression profiling for non-invasive surveillance after heart transplantation. *J Heart Lung Transpl*. 2020;39(4):S71. doi:10.1016/j.healun.2020.01.1281
68. Jordan SC, Bunnapradist S, Bromberg JS, et al. Donor-derived Cell-free DNA Identifies Antibody-mediated Rejection in Donor Specific Antibody Positive Kidney Transplant Recipients. *Transplant Direct*. 2018 Aug 20;4(9):e379.
69. CareDx. Allosure Kidney. 2021; <https://go.caredx.com/allosure/>.
70. Pai A, Swan JT, Wojciechowski D, et al. Clinical Rationale for a Routine Testing Schedule Using Donor-Derived Cell-Free DNA After Kidney Transplantation. *Ann Transplant*. Jul 02 2021; 26: e932249. PMID 34210952
71. Bloom RD, Bromberg JS, Poggio ED, et al. Cell-Free DNA and Active Rejection in Kidney Allografts. *J Am Soc Nephrol*. Jul 2017; 28(7): 2221-2232. PMID 28280140
72. Huang E, Sethi S, Peng A, et al. Early clinical experience using donor-derived cell-free DNA to detect rejection in kidney transplant recipients. *Am J Transplant*. Jun 2019; 19(6): 1663-1670. PMID 30725531
73. Stites E, Kumar D, Olaitan O, et al. High levels of dd-cfDNA identify patients with TCMR 1A and borderline allograft rejection at elevated risk of graft injury. *Am J Transplant*. Sep 2020; 20(9): 2491-2498. PMID 32056331
74. Sawinski DL, Mehta S, Alhamad T, et al. Association between dd-cfDNA levels, de novo donor specific antibodies, and eGFR decline: An analysis of the DART cohort. *Clin Transplant*. Sep 2021; 35(9): e14402. PMID 34184326

75. Mayer KA, Doberer K, Tillgren A, et al. Diagnostic value of donor-derived cell-free DNA to predict antibody-mediated rejection in donor-specific antibody-positive renal allograft recipients. *Transpl Int*. Sep 2021; 34(9): 1689-1702. PMID 34448270
76. Puliyaanda DP, Swinford R, Pizzo H, et al. Donor-derived cell-free DNA (dd-cfDNA) for detection of allograft rejection in pediatric kidney transplants. *Pediatr Transplant*. Mar 2021; 25(2): e13850. PMID 33217125
77. Sigdel TK, Archila FA, Constantin T, et al. Optimizing Detection of Kidney Transplant Injury by Assessment of Donor-Derived Cell-Free DNA via Massively Multiplex PCR. *J Clin Med*. Dec 23 2018; 8(1). PMID 30583588
78. Dengu, F. Next-generation sequencing methods to detect donor-derived cell-free DNA after transplantation, *Transplantation Reviews*, 2020 Jul;34(3):100542. doi: 10.1016/j.trre.2020.100542. Epub 2020 Mar 16.
79. Bunnapradist S, Homkrais P, Ahmed E, et al. Using both the Fraction and Quantity of Donor-Derived Cell-Free DNA to Detect Kidney Allograft Rejection. *J Am Soc Nephrol*. Oct 2021; 32(10): 2439-2441. PMID 34162734
80. Halloran PF, Reeve J, Madill-Thomsen KS, et al. Combining Donor-derived Cell-free DNA Fraction and Quantity to Detect Kidney Transplant Rejection Using Molecular Diagnoses and Histology as Confirmation. *Transplantation*. Jun 29 2022. PMID 35765145
81. Halloran PF, Reeve J, Madill-Thomsen KS, et al. The Trifecta Study: Comparing Plasma Levels of Donor-derived Cell-Free DNA with the Molecular Phenotype of Kidney Transplant Biopsies. *J Am Soc Nephrol*. Feb 2022; 33(2): 387-400. PMID 3505835477
82. Richmond ME, Deshpande SR, Zangwill SD, et al. Validation of donor fraction cell-free DNA with biopsy-proven cardiac allograft rejection in children and adults. *J Thorac Cardiovasc Surg*. Feb 2023; 165(2): 460-468.e2. PMID 35643770
83. Rodgers N, Gerding B, Cusi V, et al. Comparison of two donor-derived cell-free DNA tests and a blood gene-expression profile test in heart transplantation. *Clin Transplant*. Jun 2023; 37(6): e14984. PMID 37036133
84. Wijtvliet VPWM, Plaeke P, Abrams S, et al. Donor-derived cell-free DNA as a biomarker for rejection after kidney transplantation: a systematic review and meta-analysis. *Transpl Int*. Dec 2020; 33(12): 1626-1642. PMID 32981117
85. Xiao H, Gao F, Pang Q, et al. Diagnostic Accuracy of Donor-derived Cell-free DNA in Renal-allograft Rejection: A Meta-analysis. *Transplantation*. Jun 01 2021; 105(6): 1303-1310. PMID 32890130
86. Dandamudi R, Gu H, Goss CW, et al. Longitudinal Evaluation of Donor-Derived Cell free DNA in Pediatric Kidney Transplantation. *Clin J Am Soc Nephrol*. Nov 2022; 17(11): 1646-1655. PMID 36302566
87. Bu L, Gupta G, Pai A, et al. Clinical outcomes from the Assessing Donor-derived cell-free DNA Monitoring Insights of kidney Allografts with Longitudinal surveillance (ADMIRAL) study. *Kidney Int*. Apr 2022; 101(4): 793-803. PMID 34953773
88. Huang E, Sethi S, Peng A, et al. Early clinical experience using donor-derived cell-free DNA to detect rejection in kidney transplant recipients. *Am J Transplant*. Jun 2019; 19(6): 1663-1670. PMID 30725531
89. Sayah D, Weigt SS, Ramsey A, et al. Plasma Donor-derived Cell-free DNA Levels Are Increased During Acute Cellular Rejection After Lung Transplant: Pilot Data. *Transplant Direct*. Oct 2020; 6(10): e608. PMID 33062841
90. Khush KK, De Vlaminck I, Luikart H, et al. Donor-derived, cell-free DNA levels by next-generation targeted sequencing are elevated in allograft rejection after lung transplantation. *ERJ Open Res*. Jan 2021; 7(1). PMID 33532456

91. Keller M, Sun J, Mutebi C, et al. Donor-derived cell-free DNA as a composite marker of acute lung allograft dysfunction in clinical care. *J Heart Lung Transplant*. Apr 2022; 41(4): 458-466. PMID 35063338
92. Rosenheck JP, Ross DJ, Botros M, et al. Clinical Validation of a Plasma Donor-derived Cell-free DNA Assay to Detect Allograft Rejection and Injury in Lung Transplant. *Transplant Direct*. Apr 2022; 8(4): e1317. PMID 35372675
93. Parulekar AD, Kao CC. Detection, classification, and management of rejection after lung transplantation. *J Thorac Dis*. Sep 2019; 11(Suppl 14): S1732-S1739. PMID 31632750
94. Yusen RD, Edwards LB, Kucheryavaya AY, et al. The Registry of the International Society for Heart and Lung Transplantation: Thirty-second Official Adult Lung and Heart-Lung Transplantation Report—2015; Focus Theme: Early Graft Failure. *J Heart Lung Transplant*. Oct 2015; 34(10): 1264-77. PMID 26454740
95. Levine DJ, Glanville AR, Aboyoun C, et al. Antibody-mediated rejection of the lung: A consensus report of the International Society for Heart and Lung Transplantation. *J Heart Lung Transplant*. Apr 2016; 35(4): 397-406. PMID 27044531
96. CareDx. Allosure Lung. 2021.
97. Natera. Prospera Lung Overview. 2022.
98. Yancy CW, Jessup M, Bozkurt B, et al. 2017 ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *J Card Fail*. Aug 2017; 23(8): 628-651. PMID 28461259
99. American Society of Transplant Surgeons (ASTS). ASTS Statement on donor-derived cell-free DNA (dd-cf-DNA). 2023. Accessed on April 29, 2024. Available at: <https://asts.org/docs/default-source/position-statements/dd-cfdna-position-statement.pdf?sfvrsn=143d4bd3>
100. Velleca, et al. (2023). The International Society for Heart and Lung Transplantation (ISHLT) guidelines for the care of heart transplant recipients. *J Heart Lung Transplant*, 42(5), e1-e141. <https://doi.org/10.1016/j.healun.2022.10.015>
101. ISHLT. (2016). Antibody-mediated rejection of the lung: A consensus report of the International Society for Heart and Lung Transplantation. [https://www.jhltonline.org/article/S1053-2498\(16\)01277-8/pdf](https://www.jhltonline.org/article/S1053-2498(16)01277-8/pdf)
102. Verleden, G. M., Glanville, A. R., Lease, E. D., Fisher, A. J., Calabrese, F., Corris, P. A., Ensor, C. R., Gottlieb, J., Hachem, R. R., Lama, V., Martinu, T., Neil, D. A. H., Singer, L. G., Snell, G., & Vos, R. (2019). Chronic lung allograft dysfunction: Definition, diagnostic criteria, and approaches to treatment—A consensus report from the Pulmonary Council of the ISHLT. *J Heart Lung Transplant*, 38(5), 493-503. <https://doi.org/10.1016/j.healun.2019.03.009>
103. Kobashigawa, J., Hall, S., Shah, P., Fine, B., Halloran, P., Jackson, A. M., Khush, K. K., Margulies, K. B., Sani, M. M., Patel, J. K., Patel, N., Peyster, E., & conference, p. (2023). The evolving use of biomarkers in heart transplantation: Consensus of an expert panel. *Am J Transplant*, 23(6), 727-735. <https://doi.org/10.1016/j.ajt.2023.02.025>
104. Kasiske BL, Zeier MG, Chapman JR, et al. KDIGO clinical practice guideline for the care of kidney transplant recipients: a summary. *Kidney Int*. Feb 2010; 77(4): 299-311. PMID 19847156
105. Breda, A., Budde, K., et al. EAU Guidelines on Renal Transplantation. (2018) European Association of Urology. Accessed May 11, 2023: <https://uroweb.org/wp-content/uploads/EAU-Guidelines-on-Renal-Transplantation-2018-large-text.pdf>

106. Centers for Medicare & Medicaid Services. National Coverage Determination for HEARTSBREATH Test for Heart Transplant Rejection (260.10). 2008.
<https://www.cms.gov/medicare-coverage-database/details/ncd-details.aspx?ncdid=325&ncdver=1&keyword=heartsbreath&keywordType=starts&areald=27&docType=NCA,CAL,NCD,MEDCAC,TA,MCD,6,3,5,1,F,P&contractOption=all&sortBy=relevance&bc=AAAAAAQAAAAA&KeyWordLookUp=Doc&KeyWordSearchType=Exact>
107. Centers for Medicare and Medicaid Services. Local Coverage Determination for MoIDX: Molecular Diagnostic Tests (MDT) (L38680) original effective date 6/6/21, revision effective date 7/27/23; <https://www.cms.gov/medicare-coverage-database/view/lcd.aspx?lcdid=38680&ver=5&bc=0>.
108. Centers for Medicare and Medicaid Services. Billing and Coding: MoIDX: Molecular Testing for Solid Organ Allograft Rejection (A58207) original effective date 6/6/21, revision date 2/29/24; <https://www.cms.gov/medicare-coverage-database/view/article.aspx?articleid=58207&ver=21&bc=0>.

The articles reviewed in this research include those obtained in an Internet based literature search for relevant medical references through 4/29/24, the date the research was completed.

Joint BCBSM/BCN Medical Policy History

Policy Effective Date	BCBSM Signature Date	BCN Signature Date	Comments
1/5/06	1/5/06	1/17/06	Joint policy established
5/1/07	3/20/07	5/1/07	Maintenance – “AlloMap” added to policy
5/1/08	2/19/08	5/1/08	Routine maintenance
5/1/09	2/10/09	2/10/09	Routine maintenance
5/1/10	2/16/10	3/31/10	Policy reviewed secondary to provider inquiry and for routine maintenance. Medicare information clarified; references updated; no change to policy position. Policy title changed to “Heart Transplant Rejection Testing”. Previous title: “Laboratory Testing for Heart Transplant Rejection”
9/1/11	6/21/11	6/21/11	Routine maintenance; references updated; rationale section updated; no change to policy position.
5/1/12	2/21/12	2/21/12	Routine maintenance; references updated; no change to policy position; added code 86849 to policy.
9/1/13	6/19/13	6/26/13	Routine maintenance; references updated; no change to policy position; title changed from “Heart Transplant Rejection Testing” to current title to be consistent with BCBSA.
11/1/14	8/19/14	8/25/14	Routine maintenance; references updated; clarified coverage status for Medicare members based on CMS guidelines; no change in policy position for commercial members.
1/1/16	10/13/15	10/27/15	Routine maintenance; references updated; added code 81495, effective 1/1/16

3/1/17	12/22/16	12/27/16	<ul style="list-style-type: none"> • MPS updated added “or graft dysfunction” to Heartsbreath wording) • Position statement changed to include AlloMap with criteria • Code update (81495 deleted and replaced with 81595) • Diverges from BCBSA regarding AlloMap
3/1/18	12/12/17	12/12/17	<ul style="list-style-type: none"> • Routine maintenance
3/1/19	12/11/18		<ul style="list-style-type: none"> • Routine maintenance • Added procedure code 0055U (myTAL_{Heart} and 81479 [allosure]) • Expanded policy to address testing for kidney transplant rejection (AlloSure)
7/1/19	4/16/19		<ul style="list-style-type: none"> • Routine maintenance
7/1/20	4/14/20		<ul style="list-style-type: none"> • Routine maintenance
7/1/21	N/A		<ul style="list-style-type: none"> • Policy tabled
9/1/21	6/15/21		<ul style="list-style-type: none"> • Content and codes (83006) from Molecular Testing for CHF and Heart Transplant merged into this policy. • Title changed from Laboratory Testing for Heart and Kidney Transplant Rejection • 0118U and Viracor TRAC added • 0085T replaced with 84999 • Clarified cfDNA as EI for use post cardiac transplant (AlloSure, myTAL_{HEART}, Viracor TRAC)
9/1/22	6/21/22		<ul style="list-style-type: none"> • Routine maintenance • Added codes 0018M, 0319U, and 0320U as E/I per code update. • References updated • Evidence review for the Prospera Kidney dd-cfDNA test was added and investigational policy statement was maintained. JUMP policy aligns

			with BCBSA regarding Prospera Kidney dd-cfDNA test. Hayes has issued a D2 rating for this test.
9/1/23	6/13/23		<ul style="list-style-type: none"> • Routine maintenance • References updated • 12/15/22: Received literature review from Natera for the Prospera test. Prospera set includes the clinical validation studies for 1-Kidney Transplant monitoring (Sigdel et al and Halloran et al), 2-for use in heart transplant monitoring (Kim et al). BCBSA already incorporated these studies into the review on November 2022. • JUMP policy aligns with BCBSA regarding Prospera Kidney and Heart dd-cfDNA test as E/I. Hayes issued a D2 rating on 3/28/22 for this test. <ul style="list-style-type: none"> ○ D2 - For use of the Prospera test, which uses recipient blood to detect dd-cfDNA 1) for surveillance of rejection in post-renal-transplant patients, or 2) when clinically indicated by physician-assessed risk of active renal allograft rejection, including when a biopsy is considered or performed to evaluate suspected transplant rejection. • Vendor: N/A (ky)
9/1/24	6/14/24		<ul style="list-style-type: none"> • Routine maintenance • References updated • Incorporated IMP “Laboratory Test Post Lung Transplant for Lung Allograft Rejection – Allosure Lung”, effective July, 2023 to this JUMP policy. • Updated MPS and Exclusions section to include lung.

			<ul style="list-style-type: none"> • Title updated to Laboratory Tests Post Transplant (Kidney, Heart, and Lung) and for Heart Failure. • 4/10/24: Received and reviewed literature from Natera containing approximately 20 documents for the Prospera test requesting coverage for Heart, Lung, and Kidney. • 6/10/24: Received article 'Surveillance with dual noninvasive testing for acute cellular rejection after heart transplantation: Outcomes from the Surveillance HeartCare Outcomes Registry' published 5/15/24 from CareDx, Inc. • Vendor: N/A (ky)
--	--	--	---

Next Review Date: 2nd Qtr, 2025

BLUE CARE NETWORK BENEFIT COVERAGE
POLICY: LABORATORY TESTS POST TRANSPLANT (KIDNEY, HEART, AND LUNG) AND
FOR HEART FAILURE

I. Coverage Determination:

Commercial HMO (includes Self-Funded groups unless otherwise specified)	Covered; criteria apply
BCNA (Medicare Advantage)	Refer to the Medicare information under the Government Regulations section of this policy.
BCN65 (Medicare Complementary)	Coinsurance covered if primary Medicare covers the service.

II. Administrative Guidelines:

- The member's contract must be active at the time the service is rendered.
- Coverage is based on each member's certificate and is not guaranteed. Please consult the individual member's certificate for details. Additional information regarding coverage or benefits may also be obtained through customer or provider inquiry services at BCN.
- The service must be authorized by the member's PCP except for Self-Referral Option (SRO) members seeking Tier 2 coverage.
- Services must be performed by a BCN-contracted provider, if available, except for Self-Referral Option (SRO) members seeking Tier 2 coverage.
- Payment is based on BCN payment rules, individual certificate and certificate riders.
- Appropriate copayments will apply. Refer to certificate and applicable riders for detailed information.
- CPT - HCPCS codes are used for descriptive purposes only and are not a guarantee of coverage.